

NUCLEAR DATA AND MEASUREMENTS SERIES

ANL/NDM-65

**Note on the Elastic-Scattering of
Few-MeV Neutrons from Elemental Calcium**

by

A.B. Smith and P.T. Guenther

March 1982

**ARGONNE NATIONAL LABORATORY,
ARGONNE, ILLINOIS 60439, U.S.A.**

NUCLEAR DATA AND MEASUREMENTS SERIES


ANL/NDM-65

NOTE ON THE ELASTIC-SCATTERING OF FEW-MeV NEUTRONS
FROM ELEMENTAL CALCIUM*

by

A. B. Smith and P. T. Guenther

March 1982



U of C-AUA-USDOE

ARGONNE NATIONAL LABORATORY,
ARGONNE, ILLINOIS 60439, U.S.A.

The facilities of Argonne National Laboratory are owned by the United States Government. Under the terms of a contract (W-31-109-Eng-38) between the U. S. Department of Energy, Argonne Universities Association and The University of Chicago, the University employs the staff and operates the Laboratory in accordance with policies and programs formulated, approved and reviewed by the Association.

MEMBERS OF ARGONNE UNIVERSITIES ASSOCIATION

The University of Arizona	Kansas State University	The Ohio State University
Carnegie-Mellon University	The University of Kansas	Ohio University
Case Western Reserve University	Loyola University	The Pennsylvania State University
The University of Chicago	Marquette University	Purdue University
University of Cincinnati	Michigan State University	Saint Louis University
Illinois Institute of Technology	The University of Michigan	Southern Illinois University
University of Illinois	University of Minnesota	The University of Texas at Austin
Indiana University	University of Missouri	Washington University
Iowa State University	Northwestern University	Wayne State University
The University of Iowa	University of Notre Dame	The University of Wisconsin

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately-owned rights. Mention of commercial products, their manufacturers, or their suppliers in this publication does not imply or connote approval or disapproval of the product by Argonne National Laboratory or the U. S. Department of Energy.

ANL/NDM-65

NOTE ON THE ELASTIC-SCATTERING OF FEW-MeV NEUTRONS
FROM ELEMENTAL CALCIUM*

by

A. B. Smith and P. T. Guenther

March 1982

Applied Physics Division
Argonne National Laboratory
9700 South Cass Avenue
Argonne, Illinois 60439
U.S.A.

*This work supported by the U.S. Department of Energy.

NUCLEAR DATA AND MEASUREMENTS SERIES

The Nuclear Data and Measurements Series presents results of studies in the field of microscopic nuclear data. The primary objective is the dissemination of information in the comprehensive form required for nuclear technology applications. This Series is devoted to: a) measured microscopic nuclear parameters, b) experimental techniques and facilities employed in measurements, c) the analysis, correlation and interpretation of nuclear data, and d) the evaluation of nuclear data. Contributions to this Series are reviewed to assure technical competence and, unless otherwise stated, the contents can be formally referenced. This Series does not supplant formal journal publication but it does provide the more extensive information required for technological applications (e.g., tabulated numerical data) in a timely manner.

OTHER ISSUES IN THE ANL/NDM SERIES ARE:

- ANL/NDM-1 Cobalt Fast Neutron Cross Sections—Measurement and Evaluation by P. T. Guenther, P. A. Moldauer, A. B. Smith, D. L. Smith and J. F. Whalen, July 1973.
- ANL/NDM-2 Prompt Air-Scattering Corrections for a Fast-Neutron Fission Detector: $E_n \leq 5$ MeV by Donald L. Smith, September 1973.
- ANL/NDM-3 Neutron Scattering from Titanium; Compound and Direct Effects by E. Barnard, J. deVilliers, P. Moldauer, D. Reitmann, A. Smith and J. Whalen, October 1973.
- ANL/NDM-4 ^{90}Zr and ^{92}Zr ; Neutron Total and Scattering Cross Sections by P. Guenther, A. Smith and J. Whalen, July 1974.
- ANL/NDM-5 Delayed Neutron Data - Review and Evaluation by Samson A. Cox, April 1974.
- ANL/NDM-6 Evaluated Neutronic Cross Section File for Niobium by R. Howerton, Lawrence Livermore Laboratory and A. Smith, P. Guenther and J. Whalen, Argonne National Laboratory, May 1974.
- ANL/NDM-7 Neutron Total and Scattering Cross Sections of Some Even Isotopes of Molybdenum and the Optical Model by A. B. Smith, P. T. Guenther and J. F. Whalen, June 1974.
- ANL/NDM-8 Fast Neutron Capture and Activation Cross Sections of Niobium Isotopes by W. P. Poenitz, May 1974.
- ANL/NDM-9 Method of Neutron Activation Cross Section Measurement for $E_n = 5.5\text{--}10$ MeV Using the $\text{D}(d,n)\text{He-3}$ Reaction as a Neutron Source by D. L. Smith and J. W. Meadows, August 1974.
- ANL/NDM-10 Cross Sections for (n,p) Reactions on ^{27}Al , $^{46,47,48}\text{Tl}$, $^{54,56}\text{Fe}$, ^{58}Ni , ^{59}Co and ^{64}Zn from Near Threshold to 10 MeV by Donald L. Smith and James W. Meadows, January 1975.
- ANL/NDM-11 Measured and Evaluated Fast Neutron Cross Sections of Elemental Nickel by P. Guenther, A. Smith, D. Smith and J. Whalen, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, July 1975.
- ANL/NDM-12 A Spectrometer for the Investigation of Gamma Radiation Produced by Neutron-Induced Reactions by Donald L. Smith, April 1975.
- ANL/NDM-13 Response of Several Threshold Reactions in Reference Fission Neutron Fields by Donald L. Smith and James W. Meadows, June 1975.
- ANL/NDM-14 Cross Sections for the $^{66}\text{Zn}(n,p)^{66}\text{Cu}$, $^{113}\text{In}(n,n')^{113\text{m}}\text{In}$ and $^{115}\text{In}(n,n')^{115\text{m}}\text{In}$ Reactions from Near Threshold to 10 MeV by Donald L. Smith and James W. Meadows, July 1975.

- ANL/NDM-15 Radiative Capture of Fast Neutrons in ^{165}Ho and ^{181}Ta by W. P. Poenitz, June 1975.
- ANL/NDM-16 Fast Neutron Excitation of the Ground-State Rotational Band of ^{238}U by P. Guenther, D. Havel and A. Smith, September 1975.
- ANL/NDM-17 Sample-Size Effects in Fast-Neutron Gamma-Ray Production Measurements: Solid-Cylinder Samples by Donald L. Smith, September 1975.
- ANL/NDM-18 The Delayed Neutron Yield of ^{238}U and ^{241}Pu by J. W. Meadows January 1976.
- ANL/NDM-19 A Remark on the Prompt-Fission-Neutron Spectrum of ^{252}Cf by P. Guenther, D. Havel, R. Sjoblom and A. Smith, March 1976.
- ANL/NDM-20 Fast-Neutron-Gamma-Ray Production from Elemental Iron: $E_n \lesssim 2$ MeV by Donald L. Smith, May 1976.
- ANL/NDM-21 Note on the Experimental Determination of the Relative Fast-Neutron Sensitivity of a Hydrogenous Scintillator by A. Smith, P. Guenther and R. Sjoblom, June 1976.
- ANL/NDM-22 Note on Neutron Scattering and the Optical Model Near $A=208$ by P. Guenther, D. Havel and A. Smith, September 1976.
- ANL/NDM-23 Remarks Concerning the Accurate Measurement of Differential Cross Sections for Threshold Reactions Used in Fast-Neutron Dosimetry for Fission Reactors by Donald L. Smith, December 1976.
- ANL/NDM-24 Fast Neutron Cross Sections of Vanadium and an Evaluated Neutronic File by P. Guenther, D. Havel, R. Howerton, F. Mann, D. Smith, A. Smith and J. Whalen, May 1977.
- ANL/NDM-25 Determination of the Energy Scale for Neutron Cross Section Measurements Employing a Monoenergetic Accelerator by J. W. Meadows, January 1977.
- ANL/NDM-26 Evaluation of the $\text{IN-115}(\text{N},\text{N}')\text{IN-115M}$ Reaction for the ENDF/B-V Dosimetry File by Donald L. Smith, December 1976.
- ANL/NDM-27 Evaluated (n,p) Cross Sections of ^{46}Ti , ^{47}Ti and ^{48}Ti by C. Philis and O. Bersillon, Bruyeres-le-Chatel, France and D. Smith and A. Smith, Argonne National Laboratory, January 1977.
- ANL/NDM-28 Titanium-II: An Evaluated Nuclear Data File by C. Philis, Centre d'Etudes de Bruyères-le-Châtel, R. Howerton, Lawrence Livermore Laboratory and A. B. Smith, Argonne National Laboratory, June 1977.
- ANL/NDM-29 Note on the 250 keV Resonance in the Total Neutron Cross Section of ^6Li by A. B. Smith, P. Guenther, D. Havel and J. F. Whalen, June 1977.

- ANL/NDM-30 Analysis of the Sensitivity of Spectrum-Average Cross Sections to Individual Characteristics of Differential Excitation Functions by Donald L. Smith, March 1977.
- ANL/NDM-31 Titanium-I: Fast Neutron Cross Section Measurements by P. Guenther, D. Havel, A. Smith and J. Whalen, May 1977.
- ANL/NDM-32 Evaluated Fast Neutron Cross Section of Uranium-238 by W. Poenitz, E. Pennington, and A. B. Smith, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, October 1977.
- ANL/NDM-33 Comments on the Energy-Averaged Total Neutron Cross Sections of Structural Materials by A. B. Smith and J. F. Whalen, June 1977.
- ANL/NDM-34 Graphical Representation of Neutron Differential Cross Section Data for Reactor Dosimetry Applications by Donald L. Smith, June 1977.
- ANL/NDM-35 Evaluated Nuclear Data File of Th-232 by J. Meadows, W. Poenitz, A. Smith, D. Smith and J. Whalen, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, February 1978.
- ANL/NDM-36 Absolute Measurements of the $^{233}\text{U}(n,f)$ Cross Section Between 0.13 and 8.0 MeV by W. P. Poenitz, April 1978.
- ANL/NDM-37 Neutron Inelastic Scattering Studies for Lead-204 by D. L. Smith and J. W. Meadows, December 1977.
- ANL/NDM-38 The Alpha and Spontaneous Fission Half-Lives of ^{242}Pu by J. W. Meadows, December 1977.
- ANL/NDM-39 The Fission Cross Section of ^{239}Pu Relative to ^{235}U from 0.1 to 10 MeV by J. W. Meadows, March 1978.
- ANL/NDM-40 Statistical Theory of Neutron Nuclear Reactions by P. A. Moldauer, February 1978.
- ANL/NDM-41 Energy-Averaged Neutron Cross Sections of Fast-Reactor Structural Materials by A. Smith, R. McKnight and D. Smith, February 1978.
- ANL/NDM-42 Fast Neutron Radiative Capture Cross Section of ^{232}Th by W. P. Poenitz and D. L. Smith, March 1978.
- ANL/NDM-43 Neutron Scattering from ^{12}C in the Few-MeV Region by A. Smith, R. Holt and J. Whalen, September 1978.
- ANL/NDM-44 The Interaction of Fast Neutrons with ^{60}Ni by A. Smith, P. Guenther, D. Smith and J. Whalen, January 1979.
- ANL/NDM-45 Evaluation of $^{235}\text{U}(n,f)$ between 100 KeV and 20 MeV by W. P. Poenitz, July 1979.

- ANL/NDM-46 Fast-Neutron Total and Scattering Cross Sections of ^{107}Ag in the MeV Region by A. Smith, P. Guenther, G. Winkler and J. Whalen, January 1979.
- ANL/NDM-47 Scattering of MeV Neutrons from Elemental Iron by A. Smith and P. Guenther, March 1979.
- ANL/NDM-48 ^{235}U Fission Mass and Counting Comparison and Standardization by W. P. Poenitz, J. W. Meadows and R. J. Armani, May 1979.
- ANL/NDM-49 Some Comments on Resolution and the Analysis and Interpretation of Experimental Results from Differential Neutron Measurements by Donald L. Smith, November 1979.
- ANL/NDM-50 Prompt-Fission-Neutron Spectra of ^{233}U , ^{235}U , ^{239}Pu and ^{240}Pu Relative to that of ^{252}Cf by A. Smith, P. Guenther, G. Winkler and R. McKnight, September 1979.
- ANL/NDM-51 Measured and Evaluated Neutron Cross Sections of Elemental Bismuth by A. Smith, P. Guenther, D. Smith and J. Whalen, April 1980.
- ANL/NDM-52 Neutron Total and Scattering Cross Sections of ^6Li in the Few MeV Region by P. Guenther, A. Smith and J. Whalen, February 1980.
- ANL/NDM-53 Neutron Source Investigations in Support of the Cross Section at the Argonne Fast-Neutron Generator by James W. Meadows and Donald L. Smith, May 1980.
- ANL/NDM-54 The Nonelastic-Scattering Cross Sections of Elemental Nickel by A. B. Smith, P. T. Guenther and J. F. Whalen, June 1980.
- ANL/NDM-55 Thermal Neutron Calibration of a Tritium Extraction Facility using the $^6\text{Li}(n,t)^4\text{He}/^{197}\text{Au}(n,\gamma)^{198}\text{Au}$ Cross Section Ratio for Standardization by M. M. Bretscher and D. L. Smith, August 1980.
- ANL/NDM-56 Fast-Neutron Interactions with ^{182}W , ^{184}W and ^{186}W by P. T. Guenther, A. B. Smith and J. F. Whalen, December 1980.
- ANL/NDM-57 The Total, Elastic- and Inelastic-Scattering Fast-Neutron Cross Sections of Natural Chromium, Peter T. Guenther, Alan B. Smith and James F. Whalen, January 1981.
- ANL/NDM-58 Review of Measurement Techniques for the Neutron Capture Process by W. P. Poenitz, August 1981.
- ANL/NDM-59 Review of the Importance of the Neutron Capture Process in Fission Reactors, Wolfgang P. Poenitz, July 1981.
- ANL/NDM-60 Neutron Capture Activation Cross Sections of ^{94}Zr , ^{96}Zr , ^{98}Mo , ^{100}Mo , and ^{110}Cd , ^{114}Cd , ^{116}Cd at Thermal and 30 keV Energy, John M. Wyrick and Wolfgang P. Poenitz, (to be published).

- ANL/NDM-61 Fast-neutron Total and Scattering Cross Sections of ^{58}Ni by Carl Budtz-Jørgensen, Peter T. Guenther, Alan B. Smith and James F. Whalen, September 1981.
- ANL/NDM-62 Covariance Matrices and Applications to the Field of Nuclear Data, by Donald L. Smith, November 1981.
- ANL/NDM-63 On Neutron Inelastic-Scattering Cross Sections of ^{232}Th , ^{233}U , ^{235}U , ^{238}U , ^{239}U , and ^{239}Pu and ^{240}Pu by Alan B. Smith and Peter T. Guenther, January 1982.
- ANL/NDM-64 The Fission Fragment Angular Distributions and Total Kinetic Energies for $^{235}\text{U}(n,f)$ from .18 to 8.83 Mev by J. W. Meadows, and Carl Budtz-Jørgensen, (to be published).

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	viii
I. INTRODUCTION	1
II. EXPERIMENTAL MEASUREMENTS AND RESULTS	1
III. INTERPRETATION AND COMMENT	2

NOTE ON THE ELASTIC-SCATTERING OF FEW-MeV NEUTRONS
FROM ELEMENTAL CALCIUM

by

A. B. Smith and P. T. Guenther
Argonne National Laboratory
Argonne, Illinois 60439

ABSTRACT

Neutron differential-elastic-scattering cross sections of elemental calcium are measured from < 1.5 to 4.0 MeV at intervals of ≈ 50 keV. Scattering angles are distributed between 20 and 160 deg. Incident-neutron energy resolutions are ≈ 50 to 100 keV. The experimental results are compared with values given in ENDF/B-V and are examined in the context of shielding applications. An optical potential is deduced from the measured values and its possible implications are discussed.

*This work supported by the U.S. Department of Energy.

I. INTRODUCTION

The present experimental study had two objectives: 1) an examination of the optical potential in a region of large compound-elastic scattering (CE), double shell closure and essentially zero iso-vector component, and 2) the provision of angular-distribution information useful in shielding applications. The scattering targets were elemental calcium metal which, for the present purposes, can be considered entirely the doubly-magic nucleus ^{40}Ca . Conventional optical potentials generally include an iso-vector term^{1,2} and the imaginary portion of the potentials tends to become small in the region of shell closure.^{3,4} Quantitative knowledge of these trends is limited in the few-MeV region generally accessible only to neutron-induced processes. Calcium is a primary constituent of concrete and thus of interest from the point of view of shielding applications. First-order neutron transmission depends only upon the neutron total cross section which is reasonably well known.⁵ However, higher-order effects influencing both penetration and reflection are dependent on the scattered-neutron angular distributions. For calcium, these distributions are essentially due to elastic scattering over the several-MeV region. Elastic-scattering cross sections of calcium are known to sharply fluctuate in magnitude and angular dependence with energy throughout the few-MeV region. Previously reported measurements of calcium elastic scattering at these energies tend to be limited to isolated energies. The present measurements were so arranged as to assure continuous energy coverage with an intermediate resolution. These measurements yield a reliable knowledge of the elastic scattering to an intermediate resolution which can be further averaged to even broader resolutions to provide cross sections consistent with the concept of the optical model (OM).

II. EXPERIMENTAL MEASUREMENTS AND RESULTS

All the measurements were made using the Argonne National Laboratory ten-angle time-of-flight apparatus. That apparatus and its application have been described extensively elsewhere and will not be discussed further here.⁶ The measurements were made at incident-neutron-energy intervals of ≈ 50 keV from < 1.5 to 4.0 MeV with incident-energy resolutions of ≈ 50 to 100 keV. The scattered-neutron resolutions were sufficient to resolve the elastic-scattering component from all known inelastic-scattering contributions. Ten differential values were obtained at each incident energy, distributed between 20 and 160 deg. The scattering sample was a 2 cm in diameter and 2 cm long cylinder of calcium metal placed in a stainless-steel can 0.013 mm thick. Identical cans were available for background determinations. The statistical uncertainties of the individual measured values are $\lesssim 3\%$. The scattering angles are known to ± 1 deg. The absolute calibration of the efficiencies of the ten detectors was based upon the total cross section of elemental carbon⁷ using the methods described in Ref. 8. The detector efficiencies appeared to be known to $\pm 3\%$ and were reproducible to that accuracy. Correction procedures, such as those associated with multiple events, introduced an additional $\lesssim 1\%$ uncertainty. Thus, the overall differential-cross-section uncertainty was $\lesssim 5\%$.

The experimental results are summarized in Fig. 1. Strong energy-dependent fluctuations in both magnitude and relative shape are evident. These were smoothed by constructing a 250-keV running average of the measured values to obtain the results shown in Fig. 2. Even with the broad average, the fluctuations persist. The averaged differential cross sections were least-squared fitted with a 6th-order Legendre-polynomial series in order to obtain the angle-integrated elastic-scattering cross sections. The latter are compared in Fig. 3 with the high-resolution elastic scattering values as given in ENDF/B-V.⁹ The ENDF/B values very likely overemphasize the fluctuations above the (n;p) and (n; α) thresholds, but generally the present angle-integrated results follow the trends of the high-resolution evaluated data again with persistent fluctuations in the 250-keV averaged quantities. The same fluctuations are evident in the B_l coefficients of the Legendre-polynomial representation of the averaged values, as shown in Fig. 4, and extend through at least the B_4 term.

There have been previous measurements of calcium elastic scattering at comparable energies.^{10,11} Generally, the previous results were obtained at isolated energies which makes quantitative comparisons with the present values difficult, or even deceptive, due to the evident fluctuating structure. For example, Reber and Brandenberger¹¹ have reported results at 2.06 and 3.29 MeV. These two distributions are compared with those obtained in the present work in Fig. 5. At 2.06 MeV, the agreement is not good while at 3.29 MeV it is excellent. This dichotomy points up the importance of energy-comprehensive measurements when dealing with an energy-fluctuating cross section.

III. INTERPRETATION AND COMMENT

The thresholds for (n;p) and (n; α) reactions in calcium are at low energies and the cross sections rise rapidly to a cumulative value of ≈ 0.6 b at 4.0 MeV. Thus, one expects significant compound-nucleus (CN) decay into the charged-particle-emission channels with a corresponding dilution of the compound-elastic (CE) contribution. This eventuality complicates the interpretation of the present results. In addition, obvious large fluctuations of the elastic-scattering cross section over the energy range of the present measurements make it difficult to assure an energy-averaged behavior consistent with the concept of the optical model (OM). Faced with these problems, the OM interpretation of the experimental results was based upon the energy range ≈ 1.5 to 2.6 MeV. It was hoped that this interval was wide enough to provide a reasonable energy average in a region where the (n;p) and (n; α) contributions are yet quite small. The cumulative total of the latter is ≈ 0.1 b at 2.6 MeV and decreases rapidly with decreasing energy. Radiative neutron capture is a negligible perturbation in the present energy range and was ignored.

Calcium is a doubly-magic nucleus and it is thus appropriate to base the interpretation upon a simple spherical OM. CE contributions were large throughout the measured energy range. They were calculated using the statistical model of Hauser-Feshbach,¹² as corrected by Moldauer.¹³ All




Fig. 1. Differential-elastic-scattering cross sections of calcium. The present experimental values are denoted by data symbols. The curves result from fitting Legendre-polynomial series to the measured values. The dimensionality is scattering angle in deg. and cross section in b/sr expressed in the laboratory system.




Fig. 2. Differential-elastic-scattering cross sections of calcium. The present experimental results, averaged over 250 keV, are noted by data symbols. Curves indicate the results of model calculations as discussed in the text. The dimensionality is scattering angle in deg. and cross section in b/sr expressed in the laboratory system.




Fig. 3. Angle-integrated elastic-scattering cross sections of calcium. The present experimental results are indicated by \blacksquare . The fluctuating curve denotes the evaluated cross sections of Ref. 9. The heavy smooth curve indicates the results of model calculations as discussed in the text.




Fig. 4. Calcium B_0 to B_6 coefficients derived from the 250 keV average of the present experimental results. The B-coefficient dimensionality is barns/sr expressed in the center-of-mass system.




Fig. 5. Comparison of measured differential-elastic-scattering cross sections of calcium at 2.06 MeV (lower) and 3.29 MeV (upper). The present results are denoted by data symbols and the plain curves. Those from Ref. 11 by the curves with "tick" marks. The dimensionality is scattering angle in degrees and cross section in b/sr expressed in the center-of-mass system.

calculations utilized the spherical OM code ABAREX.¹⁴ The OM parameters were deduced by concurrently chi-square fitting all observed differential cross sections over the energy range ≈ 1.5 to 2.6 MeV. The fitting procedure simultaneously adjusted six OM parameters; real and imaginary strengths, radii, and diffusenesses. The energy dependence of the potential was taken from the global parameter set of Ref. 1. That energy dependence is of little note in the present interpretation, but it did provide for a reasonable extrapolation over a much wider energy range. The fitting procedure was pursued in a routine manner and resulted in the parameters of Table 1. Some of these parameters are unusual. The real strength (measured as Vr^2 or in terms of the integral per nucleon, J/A) is larger than conventionally encountered in "global" parameter sets.^{1,2,10} The variation can not be attributed to the iso-vector portion of the potential as that term is identically zero. Conversely, the imaginary strength (measured in terms of Wa or J/A) is unusually small. This is not so surprising as it has long been suggested that the imaginary potential is small in the region of shell closure.^{3,4} The present interpretation should be sensitive to the imaginary term as the CE component is large throughout the measured energy range.

The potential of Table 1 provides a good description of the present differential and angle-integrated results up to the onset of significant charged-particle emission as illustrated in Figs. 2 and 3. The differences between measurement and calculation are of a magnitude that might easily result from the evident fluctuations even in the illustrated 250 keV averages. Furthermore, the angle-integrated values very nicely extrapolate to low energies and are consistent with very high-energy (e.g., ≈ 10 MeV) measured values where the CE contribution is essentially negligible.^{1,11} As the (n;p) and (n; α) cross sections rapidly rise above ≈ 2.6 MeV, the calculated angle-integrated elastic scattering cross sections become consistently larger than the experimentally based quantities (extending to 6 MeV and above). This is to be expected as the decay into the charged-particle channels should significantly dilute the CE component relative to the simple model predictions which did not include charged-particle effects. Comparisons of measured and calculated differential and angle-integrated cross sections in the 3.5 to 4.0 MeV region suggests that the CN decay into the charged-particle channels amounts to 300 to 400 mb. This value is several hundred mb smaller than the cumulative (n;p) and (n; α) cross sections. Concurrently, the calculated neutron total cross section tends to be several hundred mb smaller than the measured values. These observations suggest that a significant portion of the charged-particle cross sections comes from other than CN decay. A direct charge-exchange process provides such an alternate avenue and would qualitatively augment the calculated neutron total cross section while, at the same time, reduce the charged-particle competition with the CE process. Proton and alpha-particle emission at 5.85 MeV has been studied by Foroughi and Rossel.¹⁵ These authors report a dominance of the (n;p) process with more than half of it going through the p_0 and p_1 decay branches (a cumulative cross section of 365 mb). Moreover,¹ the angular distribution of the emitted protons ($p_0 + p_1$) was observed to be very largely in the forward hemisphere in contrast to what one would expect from CN decay.

Significant fractions of the alpha-particle angular distributions were also inconsistent with the CN mechanism. Thus, at energies only somewhat above those of the present measurements, there is experimental evidence for a direct charge-exchange process which is qualitatively consistent with the above noted differences between measured and calculated differential and total cross sections. The apparent situation is unfortunate as calcium, the heaviest naturally-occurring nuclide with an identically zero iso-vector potential, is an attractive anchor point for global OM interpretations. The latter generally do not consider the direct process and, perhaps for that reason, have been troubled in their quantitative representation of high-energy calcium differential-elastic-scattering data. Evidence based upon the present work is indirect and hampered by the fluctuating nature of the cross sections at relatively low energies. However, there would have to be a relatively large deviation of the present experimental results from the true average values to appreciably influence the above conclusions.

The present experimental results improve the definition of elastic-scattering angular distributions relative to those given in ENDF/B-V. It is of interest to assess the effect of this improved definition on shielding applications. For this purpose, a simple infinite-slab Monte-Carlo calculation was carried out. The slab was assumed to consist entirely of calcium and was varied in thickness so as to provide neutron transmissions in the range ≈ 0.01 to 0.9 . The penetration through and reflection from the slab of a perpendicularly-incident neutron beam randomly distributed in energy between 1.5 and 4.0 MeV was calculated. Two data bases were employed in the calculation: i) ENDF/B-V, and ii) ENDF/B-V modified to include the additional elastic-scattering detail provided by the present measurements. The penetration and albedo of the slab calculated using the two data bases were very similar (differing by less than 1%) for all trial slab thicknesses. The differences were calculationaly significant but negligible with respect to shielding applications, particularly since any practical shield will not be limited to the pure calcium of the idealized test case. Thus, it was concluded that the improved definition of the intermediate-resolution results of the present measurements would very likely have no substantive effect on shielding considerations. This conclusion might be different for much better-resolution data.

REFERENCES

1. J. Rapaport, V. Kulkarni and R. Finlay, Nucl. Phys., A330 15(1979).
2. G. Greenlees, W. Makofske and G. Pyle, Phys. Rev., C1 1145(1970); see also Phys. Rev., C2 1063(1970).
3. W. Vonach, A. Smith and P. Moldauer, Phys. Letters, 11 331 (1964).
4. A. Lane, J. Lynn, E. Melkonian, and E. Rae, Phys. Rev. Letters, 2 424(1959).
5. See for example, Neutron Cross Sections, Vol. II, Brookhaven Nat'l. Lab. Report, BNL-325, 3rd Edition, Editors D. Garber and R. Kinsey (1976).
6. P. Guenther, A. Smith and J. Whalen, Argonne Nat'l. Lab. Report, ANL/NDM-56 (1980); see also Ref. 8.
7. A. Smith, R. Holt and J. Whalen, Argonne Nat'l Lab. Report, ANL/NDM-43 (1978); see also Ref. 9.
8. A. Smith and P. Guenther, Argonne Nat'l. Lab. Report, ANL/NDM-63 (1982).
9. Evaluated Nuclear Data File-B, Version V, Brookhaven Nat'l Lab. Report, ENDF-201 (1979), compiled by R. Kinsey.
10. B. Holmqvist and T. Wiedling, Aktiebolaget Atomenergi Report, AE-430 (1971).
11. J. Reber and J. Brandenberger, Phys. Rev., 163 1077 (1967).
12. W. Hauser and H. Feshbach, Phys. Rev., 87 366 (1952).
13. P. Moldauer, Phys. Rev., C11 426(1978); also private communication (1982).
14. ABAREX, a spherical optical-model code, P. Moldauer, private communication (1982).
15. F. Foroughi and J. Rossel, Helv. Phys. Acta, 45 439 (1972).

Table 1. Derived Optical-Potential Parameters

Real Potential ^{a,b}	
$V_0 = 56.15$	MeV
$r^c = 1.249$	F
$a = 0.491$	F
$V_0 r^2 = 87.59$	MeV • F ²
$J/A = 518.11$	MeV • F ³
$\langle r^2 \rangle = 14.28$	F ²
Imaginary Potential ^d	
$W_0 = 5.709$	MeV
$r^c = 1.328$	F
$a = 0.283$	F
$W_0 a = 1.62$	MeV • F
$J/A = 37.56$	MeV • F ³

^a Woods-Saxon form assuming $V = V_0 - 0.3 \times E$ (MeV).

^b Assuming a spin-orbit term of the Thomas form with a strength of 6.7 MeV.

^c Assuming $R = r \times A^{1/3}$.

^d Woods-Saxon derivative form assuming $W = W_0 + 0.4 \times E$ (MeV).