

NUCLEAR DATA AND MEASUREMENTS SERIES

ANL/NDM-73

**Fast-Neutron Elastic-Scattering Cross Sections
of Elemental Tin**

by

C. Budtz-Joergensen, P. Guenther, and A. Smith

July 1982

**ARGONNE NATIONAL LABORATORY,
ARGONNE, ILLINOIS 60439, U.S.A.**

NUCLEAR DATA AND MEASUREMENTS SERIES


ANL/NDM-73

Fast-Neutron Elastic-Scattering Cross
Sections of Elemental Tin*

by

C. Budtz-Jørgensen,^a P. Guenther
and A. Smith

July 1982



U of C-AUA-USDOE

ARGONNE NATIONAL LABORATORY,
ARGONNE, ILLINOIS 60439, U.S.A.

The facilities of Argonne National Laboratory are owned by the United States Government. Under the terms of a contract (W-31-109-Eng-38) among the U. S. Department of Energy, Argonne Universities Association and The University of Chicago, the University employs the staff and operates the Laboratory in accordance with policies and programs formulated, approved and reviewed by the Association.

MEMBERS OF ARGONNE UNIVERSITIES ASSOCIATION

The University of Arizona	The University of Kansas	The Ohio State University
Carnegie-Mellon University	Kansas State University	Ohio University
Case Western Reserve University	Loyola University of Chicago	The Pennsylvania State University
The University of Chicago	Marquette University	Purdue University
University of Cincinnati	The University of Michigan	Saint Louis University
Illinois Institute of Technology	Michigan State University	Southern Illinois University
University of Illinois	University of Minnesota	The University of Texas at Austin
Indiana University	University of Missouri	Washington University
The University of Iowa	Northwestern University	Wayne State University
Iowa State University	University of Notre Dame	The University of Wisconsin-Madison

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately-owned rights. Mention of commercial products, their manufacturers, or their suppliers in this publication does not imply or connote approval or disapproval of the product by Argonne National Laboratory or the U. S. Department of Energy.

ANL/NDM-73

Fast-Neutron Elastic-Scattering Cross
Sections of Elemental Tin*

by

C. Budtz-Jørgensen,^a P. Guenther
and A. Smith

July 1982

Applied Physics Division
Argonne National Laboratory
9700 South Cass Avenue
Argonne, Illinois 60439
USA

*This work supported by the U. S. Department of Energy.

^aVisiting Scientist from the Central Bureau for
Nuclear Measurements, Geel, Belgium.

NUCLEAR DATA AND MEASUREMENTS SERIES

The Nuclear Data and Measurements Series presents results of studies in the field of microscopic nuclear data. The primary objective is the dissemination of information in the comprehensive form required for nuclear technology applications. This Series is devoted to: a) measured microscopic nuclear parameters, b) experimental techniques and facilities employed in measurements, c) the analysis, correlation and interpretation of nuclear data, and d) the evaluation of nuclear data. Contributions to this Series are reviewed to assure technical competence and, unless otherwise stated, the contents can be formally referenced. This Series does not supplant formal journal publication but it does provide the more extensive information required for technological applications (e.g., tabulated numerical data) in a timely manner.

OTHER ISSUES IN THE ANL/NDM SERIES ARE:

- ANL/NDM-1 Cobalt Fast Neutron Cross Sections-Measurement and Evaluation by P. T. Guenther, P. A. Moldauer, A. B. Smith, D. L. Smith and J. F. Whalen, July 1973.
- ANL/NDM-2 Prompt Air-Scattering Corrections for a Fast-Neutron Fission Detector: $E_n \leq 5$ MeV by Donald L. Smith, September 1973.
- ANL/NDM-3 Neutron Scattering from Titanium; Compound and Direct Effects by E. Barnard, J. deVilliers, P. Moldauer, D. Reitmann, A. Smith and J. Whalen, October 1973.
- ANL/NDM-4 ^{90}Zr and ^{92}Zr ; Neutron Total and Scattering Cross Sections by P. Guenther, A. Smith and J. Whalen, July 1974.
- ANL/NDM-5 Delayed Neutron Data - Review and Evaluation by Samson A. Cox, April 1974.
- ANL/NDM-6 Evaluated Neutronic Cross Section File for Niobium by R. Howerton, Lawrence Livermore Laboratory and A. Smith, P. Guenther and J. Whalen, Argonne National Laboratory, May 1974.
- ANL/NDM-7 Neutron Total and Scattering Cross Sections of Some Even Isotopes of Molybdenum and the Optical Model by A. B. Smith, P. T. Guenther and J. F. Whalen, June 1974.
- ANL/NDM-8 Fast Neutron Capture and Activation Cross Sections of Niobium Isotopes by W. P. Poenitz, May 1974.
- ANL/NDM-9 Method of Neutron Activation Cross Section Measurement for $E_n = 5.5-10$ MeV Using the $\text{D}(d,n)\text{He-3}$ Reaction as a Neutron Source by D. L. Smith and J. W. Meadows, August 1974.
- ANL/NDM-10 Cross Sections for (n,p) Reactions on ^{27}Al , $^{46,47,48}\text{Ti}$, $^{54,56}\text{Fe}$, ^{58}Ni , ^{59}Co and ^{64}Zn from Near Threshold to 10 MeV by Donald L. Smith and James W. Meadows, January 1975.
- ANL/NDM-11 Measured and Evaluated Fast Neutron Cross Sections of Elemental Nickel by P. Guenther, A. Smith, D. Smith and J. Whalen, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, July 1975.
- ANL/NDM-12 A Spectrometer for the Investigation of Gamma Radiation Produced by Neutron-Induced Reactions by Donald L. Smith, April 1975.
- ANL/NDM-13 Response of Several Threshold Reactions in Reference Fission Neutron Fields by Donald L. Smith and James W. Meadows, June 1975.
- ANL/NDM-14 Cross Sections for the $^{66}\text{Zn}(n,p)^{66}\text{Cu}$, $^{113}\text{In}(n,n')^{113\text{m}}\text{In}$ and $^{115}\text{In}(n,n')^{115\text{m}}\text{In}$ Reactions from Near Threshold to 10 MeV by Donald L. Smith and James W. Meadows, July 1975.

- ANL/NDM-15 Radiative Capture of Fast Neutrons in ^{165}Ho and ^{181}Ta by W. P. Poenitz, June 1975.
- ANL/NDM-16 Fast Neutron Excitation of the Ground-State Rotational Band of ^{238}U by P. Guenther, D. Havel and A. Smith, September 1975.
- ANL/NDM-17 Sample-Size Effects in Fast-Neutron Gamma-Ray Production Measurements: Solid-Cylinder Samples by Donald L. Smith, September 1975.
- ANL/NDM-18 The Delayed Neutron Yield of ^{238}U and ^{241}Pu by J. W. Meadows January 1976.
- ANL/NDM-19 A Remark on the Prompt-Fission-Neutron Spectrum of ^{252}Cf by P. Guenther, D. Havel, R. Sjoblom and A. Smith, March 1976.
- ANL/NDM-20 Fast-Neutron-Gamma-Ray Production from Elemental Iron: $E_n \leq 2$ MeV by Donald L. Smith, May 1976.
- ANL/NDM-21 Note on the Experimental Determination of the Relative Fast-Neutron Sensitivity of a Hydrogenous Scintillator by A. Smith, P. Guenther and R. Sjoblom, June 1976.
- ANL/NDM-22 Note on Neutron Scattering and the Optical Model Near $A=208$ by P. Guenther, D. Havel and A. Smith, September 1976.
- ANL/NDM-23 Remarks Concerning the Accurate Measurement of Differential Cross Sections for Threshold Reactions Used in Fast-Neutron Dosimetry for Fission Reactors by Donald L. Smith, December 1976.
- ANL/NDM-24 Fast Neutron Cross Sections of Vanadium and an Evaluated Neutronic File by P. Guenther, D. Havel, R. Howerton, F. Mann, D. Smith, A. Smith and J. Whalen, May 1977.
- ANL/NDM-25 Determination of the Energy Scale for Neutron Cross Section Measurements Employing a Monoenergetic Accelerator by J. W. Meadows, January 1977.
- ANL/NDM-26 Evaluation of the $^{115}\text{In}(n,n')^{115\text{m}}\text{In}$ Reaction for the ENDF/B-V Dosimetry File by Donald L. Smith, December 1976.
- ANL/NDM-27 Evaluated (n,p) Cross Sections of ^{46}Ti , ^{47}Ti and ^{48}Ti by C. Philis and O. Bersillon, Bruyères-le-Châtel, France and D. Smith and A. Smith, Argonne National Laboratory, January 1977.
- ANL/NDM-28 Titanium-II: An Evaluated Nuclear Data File by C. Philis, Centre d'Etudes de Bruyères-le-Châtel, R. Howerton, Lawrence Livermore Laboratory and A. B. Smith, Argonne National Laboratory, June 1977.
- ANL/NDM-29 Note on the 250 keV Resonance in the Total Neutron Cross Section of ^6Li by A. B. Smith, P. Guenther, D. Havel and J. F. Whalen, June 1977.

- ANL/NDM-30 Analysis of the Sensitivity of Spectrum-Average Cross Sections to Individual Characteristics of Differential Excitation Functions by Donald L. Smith, March 1977.
- ANL/NDM-31 Titanium-I: Fast Neutron Cross Section Measurements by P. Guenther, D. Havel, A. Smith and J. Whalen, May 1977.
- ANL/NDM-32 Evaluated Fast Neutron Cross Section of Uranium-238 by W. Poenitz, E. Pennington, and A. B. Smith, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, October 1977.
- ANL/NDM-33 Comments on the Energy-Averaged Total Neutron Cross Sections of Structural Materials by A. B. Smith and J. F. Whalen, June 1977.
- ANL/NDM-34 Graphical Representation of Neutron Differential Cross Section Data for Reactor Dosimetry Applications by Donald L. Smith, June 1977.
- ANL/NDM-35 Evaluated Nuclear Data File of ^{232}Th by J. Meadows, W. Poenitz, A. Smith, D. Smith and J. Whalen, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, February 1978.
- ANL/NDM-36 Absolute Measurements of the $^{233}\text{U}(n,f)$ Cross Section Between 0.13 and 8.0 MeV by W. P. Poenitz, April 1978.
- ANL/NDM-37 Neutron Inelastic Scattering Studies for Lead-204 by D. L. Smith and J. W. Meadows, December 1977.
- ANL/NDM-38 The Alpha and Spontaneous Fission Half-Lives of ^{242}Pu by J. W. Meadows, December 1977.
- ANL/NDM-39 The Fission Cross Section of ^{239}Pu Relative to ^{235}U from 0.1 to 10 MeV by J. W. Meadows, March 1978.
- ANL/NDM-40 Statistical Theory of Neutron Nuclear Reactions by P. A. Moldauer, February 1978.
- ANL/NDM-41 Energy-Averaged Neutron Cross Sections of Fast-Reactor Structural Materials by A. Smith, R. McKnight and D. Smith, February 1978.
- ANL/NDM-42 Fast Neutron Radiative Capture Cross Section of ^{232}Th by W. P. Poenitz and D. L. Smith, March 1978.
- ANL/NDM-43 Neutron Scattering from ^{12}C in the Few-MeV Region by A. Smith, R. Holt and J. Whalen, September 1978.
- ANL/NDM-44 The Interaction of Fast Neutrons with ^{60}Ni by A. Smith, P. Guenther, D. Smith and J. Whalen, January 1979.
- ANL/NDM-45 Evaluation of $^{235}\text{U}(n,f)$ Between 100 KeV and 20 MeV by W. P. Poenitz, July 1979.

- ANL/NDM-46 Fast-Neutron Total and Scattering Cross Sections of ^{107}Ag in the MeV Region by A. Smith, P. Guenther, G. Winkler and J. Whalen, January 1979.
- ANL/NDM-47 Scattering of MeV Neutrons from Elemental Iron by A. Smith and P. Guenther, March 1979.
- ANL/NDM-48 ^{235}U Fission Mass and Counting Comparison and Standardization by W. P. Poenitz, J. W. Meadows and R. J. Armani, May 1979.
- ANL/NDM-49 Some Comments on Resolution and the Analysis and Interpretation of Experimental Results from Differential Neutron Measurements by Donald L. Smith, November 1979.
- ANL/NDM-50 Prompt-Fission-Neutron Spectra of ^{233}U , ^{235}U , ^{239}Pu and ^{240}Pu Relative to that of ^{252}Cf by A. Smith, P. Guenther, G. Winkler and R. McKnight, September 1979.
- ANL/NDM-51 Measured and Evaluated Neutron Cross Sections of Elemental Bismuth by A. Smith, P. Guenther, D. Smith and J. Whalen, April 1980.
- ANL/NDM-52 Neutron Total and Scattering Cross Sections of ^6Li in the Few MeV Region by P. Guenther, A. Smith and J. Whalen, February 1980.
- ANL/NDM-53 Neutron Source Investigations in Support of the Cross Section Program at the Argonne Fast-Neutron Generator by James W. Meadows and Donald L. Smith, May 1980.
- ANL/NDM-54 The Nonelastic-Scattering Cross Sections of Elemental Nickel by A. B. Smith, P. T. Guenther and J. F. Whalen, June 1980.
- ANL/NDM-55 Thermal Neutron Calibration of a Tritium Extraction Facility using the $^6\text{Li}(n,t)^4\text{He}/^{197}\text{Au}(n,\gamma)^{198}\text{Au}$ Cross Section Ratio for Standardization by M. M. Bretscher and D. L. Smith, August 1980.
- ANL/NDM-56 Fast-Neutron Interactions with ^{182}W , ^{184}W and ^{186}W by P. T. Guenther, A. B. Smith and J. F. Whalen, December 1980.
- ANL/NDM-57 The Total, Elastic- and Inelastic-Scattering Fast-Neutron Cross Sections of Natural Chromium, Peter T. Guenther, Alan B. Smith and James F. Whalen, January 1981.
- ANL/NDM-58 Review of Measurement Techniques for the Neutron Capture Process by W. P. Poenitz, August 1981.
- ANL/NDM-59 Review of the Importance of the Neutron Capture Process in Fission Reactors, Wolfgang P. Poenitz, July 1981.
- ANL/NDM-60 Neutron Capture Activation Cross Sections of ^{94}Zr , ^{96}Zr , $^{98,100}\text{Mo}$, and $^{110,114,116}\text{Cd}$ at Thermal and 30 keV Energy, John M. Wyrick and Wolfgang P. Poenitz, (to be published).

- ANL/NDM-61 Fast-neutron Total and Scattering Cross Sections of ^{58}Ni by Carl Budtz-Jørgensen, Peter T. Guenther, Alan B. Smith and James F. Whalen, September 1981.
- ANL/NDM-62 Covariance Matrices and Applications to the Field of Nuclear Data, by Donald L. Smith, November 1981.
- ANL/NDM-63 On Neutron Inelastic-Scattering Cross Sections of ^{232}Th , ^{233}U , ^{235}U , ^{238}U , ^{239}U , and ^{239}Pu and ^{240}Pu by Alan B. Smith and Peter T. Guenther, January 1982.
- ANL/NDM-64 The Fission Fragment Angular Distributions and Total Kinetic Energies for $^{235}\text{U}(n,f)$ from 0.18 to 8.83 MeV by James W. Meadows, and Carl Budtz-Jørgensen, (to be published).
- ANL/NDM-65 Note on the Elastic Scattering of Several MeV Neutrons from Elemental Calcium by Alan B. Smith and Peter T. Guenther, March 1982.
- ANL/NDM-66 Fast-neutron Scattering Cross Sections of Elemental Silver by Alan B. Smith and Peter T. Guenther, May 1982.
- ANL/NDM-67 Non-evaluation Applications for Covariance Matrices by Donald L. Smith, July 1982.
- ANL/NDM-68 Fast-neutron Total and Scattering Cross Sections of ^{103}Rh by Alan B. Smith, Peter T. Guenther and James F. Whalen (to be published).
- ANL/NDM-69 Fast-neutron Scattering Cross Sections of Elemental Zirconium by Alan B. Smith and Peter T. Guenther (to be published).
- ANL/NDM-70 Fast-neutron Total and Scattering Cross Sections of Niobium by Alan B. Smith, Peter T. Guenther and James F. Whalen (to be published).
- ANL/NDM-71 Fast-neutron Total and Scattering Cross Sections of Elemental Palladium by Alan B. Smith, Peter T. Guenther and James F. Whalen, June 1982.
- ANL/NDM-72 Fast-neutron Scattering from Elemental Cadmium by Alan B. Smith and P. T. Guenther (to be published).

Table of Contents

	<u>Page</u>
ABSTRACT	viii
I. INTRODUCTION	1
II. BRIEF OUTLINE OF EXPERIMENTAL METHODS	1
III. EXPERIMENTAL RESULTS	2
IV. MODEL DERIVATION	2
V. COMPARISONS WITH ENDF/B-V	5
VI. SUMMARY COMMENTS	8
ACKNOWLEDGEMENTS	8

Fast-Neutron Elastic-Scattering Cross
Sections of Elemental Tin

by

C. Budtz-Jørgensen, P. T. Guenther
and A. B. Smith

Applied Physics Division
Argonne National Laboratory
Argonne, Illinois 60439

ABSTRACT

Broad-resolution neutron-elastic-scattering cross sections of elemental tin are measured from 1.5 to 4.0 MeV. Incident-energy intervals are ≈ 50 keV below 3.0 MeV and ≈ 200 keV at higher energies. Ten to twenty scattering angles are used, distributed between ≈ 20 and 160° . The experimental results are used to deduce the parameters of a spherical optical-statistical model and they are also compared with corresponding values given in ENDF/B-V.

I. INTRODUCTION

The present work was undertaken as part of a comprehensive study of fast-neutron interactions with the light-mass fission products including neutron total, scattering and capture cross sections¹⁻⁷. The tin isotopes are at the upper extreme of the light-mass fission-product yield distribution. Typically, their fission-product yields are ¹¹²Sn (0.1%), ¹¹⁴Sn(0.05%), ¹¹⁵Sn(0.03%), ¹¹⁶Sn(0.044%), ¹¹⁷Sn(0.042%), ¹¹⁸Sn(0.041%), ¹¹⁹Sn(0.041%), ¹²⁰Sn(0.42%), ¹²²Sn(0.045%) and ¹²⁴Sn(0.056%) for ²³⁹Pu fission induced by thermal neutrons. These are relatively small values. However, the neutron interaction with tin is of fission-product significance as it is a basis for model parameters that are generally applicable in this mass region. The present work was undertaken in order to provide an experimental foundation and to derive therefrom a representative set of model parameters. The latter were subsequently employed in the formulation of a "regional" model applicable to the light-mass fission products⁸. Elemental tin consists of the ten isotopes ¹¹²Sn(1.0%), ¹¹⁴Sn(0.86%), ¹¹⁵Sn(0.35%), ¹¹⁶Sn(14.4%), ¹¹⁷Sn(7.6%), ¹¹⁸Sn(24.1%), ¹¹⁹Sn(8.6%), ¹²⁰Sn(32.8%), ¹²²Sn(4.7%) and ¹²⁴Sn(5.8%). These isotopes cover a large range of the nuclear asymmetry parameter, (N-Z)/A. However, the isotopic abundance is concentrated in the three isotopes ¹¹⁶Sn, ¹¹⁸Sn and ¹²⁰Sn. Thus the present elemental measurements and model are representative of the neutron interaction with targets having similar asymmetries.

Subsequent portions of this paper address; i) a brief outline of the experimental method, ii) the experimental results, iii) the derivation of an optical-statistical model (OM) from the measured values, and iv) comparisons of the experimental results with the evaluated quantities given in ENDF/B-V⁹.

II. BRIEF OUTLINE OF THE EXPERIMENTAL METHOD

The measurements were made using the time-of-flight technique with the Argonne ten-angle detection apparatus¹⁰. The neutron source was the ⁷Li(p,n)⁷Be reaction pulsed on for durations of ≈ 1 nsec at a repetition rate of 2 MHz. The scattering sample was a cylinder of elemental tin 2 cm in diameter and 2 cm long placed 13 cm from the neutron source at a zero-degree source-reaction angle. Ten proton-recoil-scintillation detectors were placed ≈ 5.4 m from the scattering sample distributed over scattering angles from 20 to 160°. The relative scattering angles were known to $\pm 0.2^\circ$ and the absolute angular scale to $\pm 0.6^\circ$. The relative detector efficiencies were determined by the observation of neutrons emitted from the spontaneous fission of ²⁵²Cf¹¹. The absolute normalization of the detector efficiencies was determined relative to the neutron total cross sections of elemental carbon¹² in the manner described in ref. 13. These procedures implied that the differential-scattering measurements were made relative to the neutron total cross sections of carbon. The experimental results were corrected for multiple-event, beam-attenuation and angular-resolution effects using a combination of Monte-Carlo and analytical procedures as described in ref. 14. Generally, details of the measurement method are set forth in ref. 14.

III. EXPERIMENTAL RESULTS

The measurements were made from 1.5 to 4.0 MeV with incident-energy spreads of 40 to 70 keV. Below 3.0 MeV the incident-energy-measurement interval was ≈ 50 keV and ten scattering angles were used. In order to reduce the effect of any possible cross-section fluctuations the results obtained at adjacent incident energies were averaged to a single distribution. At energies above 3.0 MeV the measurements were made at 200 keV incident-energy intervals and twenty scattering angles. The statistical uncertainties of the individual differential elastic-scattering measurements were $\lesssim 1\%$ excepting the minima regions of the higher-energy distributions where they were larger. The uncertainties associated with the detector normalizations were $\approx 3\%$ and those due to the correction procedures $\lesssim 1\%$. Thus, the overall uncertainties of the measured differential-elastic-scattering cross sections were $\lesssim 5\%$. The energy-averaged elastic-scattering results are summarized in Fig. 1. The corresponding numerical values have been transmitted to the National Nuclear Data Center, Brookhaven National Laboratory. The angle-integrated elastic-scattering cross sections were derived from the measured differential values by least-squares fitting the observed differential distributions with 6th-order Legendre-Polynomial series. The results of these fitting procedures were descriptive of the observations, as illustrated in Fig. 1. The corresponding angle-integrated elastic-scattering cross sections are shown in Fig. 2. Their uncertainties are $\lesssim 5\%$. The energy-averaged experimental results displayed no evidence of energy-dependent fluctuations. There have been a number of previously reported measurements of differential-elastic scattering from elemental tin as referenced in CINDA¹⁶. These prior results should be reasonably summarized by the evaluations of ENDF/B-V⁹ and comparisons with those evaluations are discussed below.

Neutrons due to inelastic-scattering processes were observed in the course of the above elastic-scattering measurements. They corresponded to relatively high excitation energies and were attributed to contributions from a number of isotopes. The primary purpose of the present measurements was the acquisition of precise energy-averaged elastic-scattering cross sections. To achieve that objective, relatively broad incident-energy spreads were used and these precluded detailed resolution of the complex inelastic-neutron components. Therefore, cross sections were not derived from the inelastic-neutron data. Detailed experimental studies of the inelastic-scattering process in tin will be reported later.

IV. MODEL DERIVATION

It was assumed that the present experimental results could be described in terms of a the spherical optical-statistical model (OM)¹⁷. Approximately




Fig. 1. Measured Differential-Elastic-Scattering Cross Sections of Elemental Tin. The present experimental results are indicated by data symbols and curves are the results of Legendre-Polynomial fits to the data.




Fig. 2. Neutron Total and Angle-Integrated Elastic-Scattering Cross Sections of Elemental Tin. The present experimental results, and lower-energy (< 1.5 MeV) elastic-scattering results of ref. 15, are noted by \circ and a 50 keV average of the neutron total cross sections of ref. 1 by \times . Curves indicate the results of calculations as described in the text.

85% of the element consists of even isotopes and that portion is primarily due to the three isotopes ^{116}Sn , ^{118}Sn and ^{120}Sn . Therefore, this OM derivation assumed an elemental mass of 118.8 AMU and the level structure of ^{118}Sn as reasonably representative of that of the element. In the energy range of the present measurements compound-nucleus processes are prominent. They were calculated using the Hauser-Feshbach formula¹⁸, as modified by Moldauer¹⁹. The excitation energies and $J-\pi$ values of ^{118}Sn were taken from ref. 20 up to excitations of 2.4 MeV. Higher-energy excitations were described in terms of the statistical formalism and parameters of Gilbert and Cameron²¹. All of the calculations were carried out using the computer code ABAREX²².

The entire elastic-scattering data base (Fig. 1) was concurrently chi-square fitted, simultaneously varying the six parameters; real and imaginary strengths, radii and diffusenesses. The real strength was assumed to have an energy dependence of the form $V = V_0 - 0.3 \cdot E$ (MeV). In addition, a 6 MeV spin-orbit potential of the Thomas form was assumed. The fitting procedure rapidly converged to the parameters of Table 1. The real parameters are similar to those reported in the literature²³. The imaginary strength is relatively small, as previously reported near shell closures²⁴ ($Z = 50$ for tin). The parameters of Table 1 give a good description of the observed differential-elastic-scattering cross sections, as illustrated in Fig. 3. More than 95% of the measured values are consistent with the calculated quantities to within the experimental uncertainties. The parameters of Table 1 provide a good description of the angle-integrated elastic-scattering cross sections and of the neutron total cross sections of ref. 1 as illustrated in Fig. 2.

The agreement between observation and calculation supports the consideration of the OM parameters of Table 1 in the formulation of a "regional" OM applicable to the light-mass fission-product region.

V. COMPARISONS WITH ENDF/B-V

The ENDF/B-V⁹ evaluated data file gives neutron cross sections of tin on an isotopic basis. These isotopic components were combined to obtain an evaluated elemental file for comparisons with the present experimental values. The results of such comparisons are summarized in Fig. 4 and Table 2. Above several MeV, the experimental neutron total cross sections of ref. 1 are consistent with those of the file. At lower energies the measured values are larger than the evaluated quantities by a few percent. This difference is reflected in comparisons of lower-energy measured and evaluated elastic-scattering cross sections as the radiative-capture cross sections are relatively small above several-100 keV. However, in the energy range of the present experiments the evaluated and measured elastic-scattering cross sections differ by large amounts: e.g. up to 20-30%. The latter discrepancies are far beyond the relevant experimental uncertainties or reasonable contributions from the low-lying levels of ^{119}Sn (8.6% abundant). In addition, the evaluated elastic-scatter-




Fig. 3. Comparisons of Measured (symbols) and Calculated (curves) Neutron Differential-Elastic-Scattering Cross Sections of Elemental Tin.




Fig. 4. Comparison of Measured and Evaluated Neutron Total and Elastic-Scattering Cross Sections of Elemental Tin. "Eyeguides", constructed through the present experimental results and those of ref. 1, are indicated by heavy curves. Light curves denote the comparable values implied by ENDF/B-V9.

ing cross sections at higher energies, extending up to 20 MeV, appear to have shapes that are not consistent with a conventional OM. The above situation results in considerable differences between measured and evaluated nonelastic cross sections amounting to more than 100% at some energies. At the energies of the present measurements the nonelastic cross sections are essentially the total inelastic-scattering cross sections. Thus the nonelastic discrepancies are equivalent to inelastic-scattering discrepancies. They are factors of up to five larger than the often cited 20% accuracy goal for fission-product inelastic-scattering cross sections.

VI. SUMMARY COMMENTS

The present experimental results give improved definition to neutron elastic-scattering from elemental tin over the energy range 1.5 to 4.0 MeV. The measured values were used to determine the parameters of a spherical optical-statistical model. The resulting parameters provided a good description of the observed differential-elastic-scattering cross sections and of recently measured neutron total cross sections¹ over the few-MeV range. The model parameters suggest a relatively weak absorption similar to that previously cited in regions of shell closure ($Z = 50$)²⁴. The measured elastic-scattering cross sections are much smaller than the comparable elemental values deduced from ENDF/B-V⁹, by amounts that considerably exceed the experimental uncertainties. The present experimental results, and the neutron total cross sections of ref. 1, imply nonelastic cross sections that are much larger than those given by the evaluation. These differences are reflected in similar large discrepancies between experimentally-implied and evaluated total-inelastic-scattering cross sections by amounts very much exceeding the frequently cited applied accuracy goals for fission-product inelastic-scattering cross sections.

ACKNOWLEDGEMENTS

The authors are indebted to Drs. W. Poenitz and P. Moldauer for helpful discussions and the provision of pre-publication data. They are also grateful to Dr. E. Pennington for assistance in handling the ENDF/B-V files.

References

1. W. Poenitz and J. Whalen, to be published; see also Nucl. Sci. and Eng., 78 333 (1981) for details of the total-cross-section measurement methods.
2. J. Wyrick and W. Poenitz, Proc. NEANDC/NEACRP Specialist's Mtg. on Fast-Neutron Capture, Argonne (1982).
3. A. B. Smith and P. T. Guenther, Argonne National Laboratory Report, ANL/NDM-66 (1982).
4. A. B. Smith, P. T. Guenther and J. Whalen, Argonne National Laboratory Report, ANL/NDM-68 (1982).
5. A. B. Smith and P. T. Guenther, Argonne National Laboratory Report, ANL/NDM-69 (1982).
6. A. B. Smith, P. Guenther and J. Whalen, Argonne National Laboratory Report, ANL/NDM-70 (1982).
7. A. B. Smith and P. T. Guenther, Argonne National Laboratory Report, ANL/NDM-71 (1982).
8. A. B. Smith, (to be published) (1982).
9. Evaluated Nuclear Data File-B, Version V, Brookhaven National Laboratory Report, ENDF-201 (1979), compiled by R. Kinsey.
10. A. B. Smith, P. T. Guenther, R. Larsen, C. Nelson, P. Walker and J. Whalen, Nucl. Instr. and Meth. 50 277 (1967).
11. A. B. Smith, P. T. Guenther and R. Sjoblom, Nucl. Instr. and Meth., 140 397 (1977).
12. A. B. Smith, P. T. Guenther, R. Holt and J. Whalen, Nucl. Sci. and Eng., 70 281 (1979).
13. A. B. Smith and P. T. Guenther, Argonne National Laboratory Report, ANL/NDM-63 (1982).
14. P. T. Guenther, Elastic and Inelastic Scattering of Fast Neutrons from the Even Isotopes of Tungsten, Univ. of Illinois Thesis (1977).
15. A. Smith and R. Hayes, Nucl. Phys., A93 609 (1967).
16. CINDA-81, An Index to the Literature on Microscopic Neutron Data, IAEA Press, Vienna (1981).
17. P. Hodgson, Nuclear Reactions and Nuclear Structure, Clarendon Press, Oxford (1971).
18. W. Hauser and H. Feshbach, Phys. Rev., 87 366 (1952).

19. P. A. Moldauer, Phys. Rev., C11 426 (1978); also private communication.
20. Table of Isotopes, 7th Edition, C. M. Lederer and V. S. Shirley. Eds., John Wiley and Sons, Inc. New York (1978).
21. A. Gilbert and A. Cameron, Can. Jour. of Phys., 43 1446 (1965).
22. ABAREX, a Spherical Optical-Model Code, P. A. Moldauer, private communication (1982).
23. J. Rapaport, V. Kulparni and R. Finley, Nucl. Phys. A330 15 (1979).
24. W. Vonach, A. B. Smith and P. T. Guenther, Phys. Lett., 11 331 (1964).

Table 1. Optical-Model Parameters for Elemental Tin

Real Potential

Strength ^a	$V_0 = 48.528$	MeV
Radius ^b	$r_v = 1.251$	F
Diffuseness	$a_v = 0.596$	F
$V_r^2 = 75.94$	MeV-F^2	
$J_v/A^d = 434.86$	MeV-F^3	

Imaginary Potential

Strength ^c	$W = 7.04$	MeV
Radius	$r_w = 1.244$	F
Diffuseness	$a_w = 0.4707$	F
$W_a = 3.31$	MeV-F	
$J_w/A = 53.49$	MeV-F^3	

-
- a. Saxon form. Assume energy dependence of the form $V = V_0 - 0.3 \times E(\text{MeV})$ and a spin-orbit potential of the Thomas form with a 6 MeV strength.
 - b. All radii expressed as $R = r_0 \times A^{1/3}$.
 - c. Saxon derivative form.
 - d. Integral per nucleon given by J/A .

Table 2. Comparisons with ENDF/B-V

E_n (MeV)	$\frac{\text{ENDF-Exp.}}{\text{ENDF}}$		
	σ_{total}	σ_{elastic}	$\sigma_{\text{nonelastic}}$
0.5	-3.5%	---	---
1.0	-3.9%	---	---
1.5	-2.2%	- 3.4%	+ 5.5%
2.0	-1.3%	0.0%	- 7.5%
2.5	-1.0%	+15.8%	- 92.1%
3.0	-0.8%	+25.3%	-129.5%
3.5	+0.7%	+27.6%	- 85.0%
4.0	+0.5%	+24.1%	- 50.4%
4.5	+0.0%	---	---