

**NUCLEAR DATA AND MEASUREMENTS SERIES**

**ANL/NDM-75**

**Neutron Total and Scattering Cross Sections  
of Elemental Antimony**

by

A.B. Smith, P.T. Guenther, and J.F. Whalen

November 1982

**ARGONNE NATIONAL LABORATORY,  
ARGONNE, ILLINOIS 60439, U.S.A.**

# NUCLEAR DATA AND MEASUREMENTS SERIES


ANL/NDM-75

## NEUTRON TOTAL AND SCATTERING CROSS SECTIONS OF ELEMENTAL ANTIMONY\*

by

A. B. Smith, P. T. Guenther and J. F. Whalen<sup>1</sup>

November 1982



U of C-AUA-USDOE

ARGONNE NATIONAL LABORATORY,  
ARGONNE, ILLINOIS 60439, U.S.A.

The facilities of Argonne National Laboratory are owned by the United States Government. Under the terms of a contract (W-31-109-Eng-38) among the U. S. Department of Energy, Argonne Universities Association and The University of Chicago, the University employs the staff and operates the Laboratory in accordance with policies and programs formulated, approved and reviewed by the Association.

#### MEMBERS OF ARGONNE UNIVERSITIES ASSOCIATION

The University of Arizona	The University of Kansas	The Ohio State University
Carnegie-Mellon University	Kansas State University	Ohio University
Case Western Reserve University	Loyola University of Chicago	The Pennsylvania State University
The University of Chicago	Marquette University	Purdue University
University of Cincinnati	The University of Michigan	Saint Louis University
Illinois Institute of Technology	Michigan State University	Southern Illinois University
University of Illinois	University of Minnesota	The University of Texas at Austin
Indiana University	University of Missouri	Washington University
The University of Iowa	Northwestern University	Wayne State University
Iowa State University	University of Notre Dame	The University of Wisconsin-Madison

#### NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately-owned rights. Mention of commercial products, their manufacturers, or their suppliers in this publication does not imply or connote approval or disapproval of the product by Argonne National Laboratory or the U. S. Department of Energy.

ANL/NDM-75

NEUTRON TOTAL AND SCATTERING  
CROSS SECTIONS OF ELEMENTAL ANTIMONY\*

by

A. B. Smith, P. T. Guenther and J. F. Whalen

November 1982

Applied Physics Division  
Argonne National Laboratory  
9700 South Cass Avenue  
Argonne, Illinois 60439  
U.S.A.

---

\*Work supported by the U.S. Department of Energy.

## NUCLEAR DATA AND MEASUREMENTS SERIES

The Nuclear Data and Measurements Series presents results of studies in the field of microscopic nuclear data. The primary objective is the dissemination of information in the comprehensive form required for nuclear technology applications. This Series is devoted to: a) measured microscopic nuclear parameters, b) experimental techniques and facilities employed in measurements, c) the analysis, correlation and interpretation of nuclear data, and d) the evaluation of nuclear data. Contributions to this Series are reviewed to assure technical competence and, unless otherwise stated, the contents can be formally referenced. This Series does not supplant formal journal publication but it does provide the more extensive information required for technological applications (e.g., tabulated numerical data) in a timely manner.

OTHER ISSUES IN THE ANL/NDM SERIES ARE:

- ANL/NDM-1 Cobalt Fast Neutron Cross Sections-Measurement and Evaluation by P. T. Guenther, P. A. Moldauer, A. B. Smith, D. L. Smith and J. F. Whalen, July 1973.
- ANL/NDM-2 Prompt Air-Scattering Corrections for a Fast-Neutron Fission Detector:  $E_n \leq 5$  MeV by Donald L. Smith, September 1973.
- ANL/NDM-3 Neutron Scattering from Titanium; Compound and Direct Effects by E. Barnard, J. deVilliers, P. Moldauer, D. Reitmann, A. Smith and J. Whalen, October 1973.
- ANL/NDM-4  $^{90}\text{Zr}$  and  $^{92}\text{Zr}$ ; Neutron Total and Scattering Cross Sections by P. Guenther, A. Smith and J. Whalen, July 1974.
- ANL/NDM-5 Delayed Neutron Data - Review and Evaluation by Samson A. Cox, April 1974.
- ANL/NDM-6 Evaluated Neutronic Cross Section File for Niobium by R. Howerton, Lawrence Livermore Laboratory and A. Smith, P. Guenther and J. Whalen, Argonne National Laboratory, May 1974.
- ANL/NDM-7 Neutron Total and Scattering Cross Sections of Some Even Isotopes of Molybdenum and the Optical Model by A. B. Smith, P. T. Guenther and J. F. Whalen, June 1974.
- ANL/NDM-8 Fast Neutron Capture and Activation Cross Sections of Niobium Isotopes by W. P. Poenitz, May 1974.
- ANL/NDM-9 Method of Neutron Activation Cross Section Measurement for  $E_n = 5.5-10$  MeV Using the  $\text{D}(d,n)\text{He-3}$  Reaction as a Neutron Source by D. L. Smith and J. W. Meadows, August 1974.
- ANL/NDM-10 Cross Sections for (n,p) Reactions on  $^{27}\text{Al}$ ,  $^{46,47,48}\text{Ti}$ ,  $^{54,56}\text{Fe}$ ,  $^{58}\text{Ni}$ ,  $^{59}\text{Co}$  and  $^{64}\text{Zn}$  from Near Threshold to 10 MeV by Donald L. Smith and James W. Meadows, January 1975.
- ANL/NDM-11 Measured and Evaluated Fast Neutron Cross Sections of Elemental Nickel by P. Guenther, A. Smith, D. Smith and J. Whalen, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, July 1975.
- ANL/NDM-12 A Spectrometer for the Investigation of Gamma Radiation Produced by Neutron-Induced Reactions by Donald L. Smith, April 1975.
- ANL/NDM-13 Response of Several Threshold Reactions in Reference Fission Neutron Fields by Donald L. Smith and James W. Meadows, June 1975.
- ANL/NDM-14 Cross Sections for the  $^{66}\text{Zn}(n,p)^{66}\text{Cu}$ ,  $^{113}\text{In}(n,n')^{113\text{m}}\text{In}$  and  $^{115}\text{In}(n,n')^{115\text{m}}\text{In}$  Reactions from Near Threshold to 10 MeV by Donald L. Smith and James W. Meadows, July 1975.

- ANL/NDM-15 Radiative Capture of Fast Neutrons in  $^{165}\text{Ho}$  and  $^{181}\text{Ta}$  by W. P. Poenitz, June 1975.
- ANL/NDM-16 Fast Neutron Excitation of the Ground-State Rotational Band of  $^{238}\text{U}$  by P. Guenther, D. Havel and A. Smith, September 1975.
- ANL/NDM-17 Sample-Size Effects in Fast-Neutron Gamma-Ray Production Measurements: Solid-Cylinder Samples by Donald L. Smith, September 1975.
- ANL/NDM-18 The Delayed Neutron Yield of  $^{238}\text{U}$  and  $^{241}\text{Pu}$  by J. W. Meadows January 1976.
- ANL/NDM-19 A Remark on the Prompt-Fission-Neutron Spectrum of  $^{252}\text{Cf}$  by P. Guenther, D. Havel, R. Sjoblom and A. Smith, March 1976.
- ANL/NDM-20 Fast-Neutron-Gamma-Ray Production from Elemental Iron:  $E_n \lesssim 2$  MeV by Donald L. Smith, May 1976.
- ANL/NDM-21 Note on the Experimental Determination of the Relative Fast-Neutron Sensitivity of a Hydrogenous Scintillator by A. Smith, P. Guenther and R. Sjoblom, June 1976.
- ANL/NDM-22 Note on Neutron Scattering and the Optical Model Near  $A=208$  by P. Guenther, D. Havel and A. Smith, September 1976.
- ANL/NDM-23 Remarks Concerning the Accurate Measurement of Differential Cross Sections for Threshold Reactions Used in Fast-Neutron Dosimetry for Fission Reactors by Donald L. Smith, December 1976.
- ANL/NDM-24 Fast Neutron Cross Sections of Vanadium and an Evaluated Neutronic File by P. Guenther, D. Havel, R. Howerton, F. Mann, D. Smith, A. Smith and J. Whalen, May 1977.
- ANL/NDM-25 Determination of the Energy Scale for Neutron Cross Section Measurements Employing a Monoenergetic Accelerator by J. W. Meadows, January 1977.
- ANL/NDM-26 Evaluation of the  $^{115}\text{In}(n,n')^{115\text{m}}\text{In}$  Reaction for the ENDF/B-V Dosimetry File by Donald L. Smith, December 1976.
- ANL/NDM-27 Evaluated (n,p) Cross Sections of  $^{46}\text{Ti}$ ,  $^{47}\text{Ti}$  and  $^{48}\text{Ti}$  by C. Philis and O. Bersillon, Bruyeres-le-Chatel, France and D. Smith and A. Smith, Argonne National Laboratory, January 1977.
- ANL/NDM-28 Titanium-II: An Evaluated Nuclear Data File by C. Philis, Centre d'Etudes de Bruyères-le-Châtel, R. Howerton, Lawrence Livermore Laboratory and A. B. Smith, Argonne National Laboratory, June 1977.
- ANL/NDM-29 Note on the 250 keV Resonance in the Total Neutron Cross Section of  $^6\text{Li}$  by A. B. Smith, P. Guenther, D. Havel and J. F. Whalen, June 1977.

- ANL/NDM-30 Analysis of the Sensitivity of Spectrum-Average Cross Sections to Individual Characteristics of Differential Excitation Functions by Donald L. Smith, March 1977.
- ANL/NDM-31 Titanium-I: Fast Neutron Cross Section Measurements by P. Guenther, D. Havel, A. Smith and J. Whalen, May 1977.
- ANL/NDM-32 Evaluated Fast Neutron Cross Section of Uranium-238 by W. Poenitz, E. Pennington, and A. B. Smith, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, October 1977.
- ANL/NDM-33 Comments on the Energy-Averaged Total Neutron Cross Sections of Structural Materials by A. B. Smith and J. F. Whalen, June 1977.
- ANL/NDM-34 Graphical Representation of Neutron Differential Cross Section Data for Reactor Dosimetry Applications by Donald L. Smith, June 1977.
- ANL/NDM-35 Evaluated Nuclear Data File of  $^{232}\text{Th}$  by J. Meadows, W. Poenitz, A. Smith, D. Smith and J. Whalen, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, February 1978.
- ANL/NDM-36 Absolute Measurements of the  $^{233}\text{U}(n,f)$  Cross Section Between 0.13 and 8.0 MeV by W. P. Poenitz, April 1978.
- ANL/NDM-37 Neutron Inelastic Scattering Studies for Lead-204 by D. L. Smith and J. W. Meadows, December 1977.
- ANL/NDM-38 The Alpha and Spontaneous Fission Half-Lives of  $^{242}\text{Pu}$  by J. W. Meadows, December 1977.
- ANL/NDM-39 The Fission Cross Section of  $^{239}\text{Pu}$  Relative to  $^{235}\text{U}$  from 0.1 to 10 MeV by J. W. Meadows, March 1978.
- ANL/NDM-40 Statistical Theory of Neutron Nuclear Reactions by P. A. Moldauer, February 1978.
- ANL/NDM-41 Energy-Averaged Neutron Cross Sections of Fast-Reactor Structural Materials by A. Smith, R. McKnight and D. Smith, February 1978.
- ANL/NDM-42 Fast Neutron Radiative Capture Cross Section of  $^{232}\text{Th}$  by W. P. Poenitz and D. L. Smith, March 1978.
- ANL/NDM-43 Neutron Scattering from  $^{12}\text{C}$  in the Few-MeV Region by A. Smith, R. Holt and J. Whalen, September 1978.
- ANL/NDM-44 The Interaction of Fast Neutrons with  $^{60}\text{Ni}$  by A. Smith, P. Guenther, D. Smith and J. Whalen, January 1979.
- ANL/NDM-45 Evaluation of  $^{235}\text{U}(n,f)$  Between 100 KeV and 20 MeV by W. P. Poenitz, July 1979.

- ANL/NDM-46 Fast-Neutron Total and Scattering Cross Sections of  $^{107}\text{Ag}$  in the MeV Region by A. Smith, P. Guenther, G. Winkler and J. Whalen, January 1979.
- ANL/NDM-47 Scattering of MeV Neutrons from Elemental Iron by A. Smith and P. Guenther, March 1979.
- ANL/NDM-48  $^{235}\text{U}$  Fission Mass and Counting Comparison and Standardization by W. P. Poenitz, J. W. Meadows and R. J. Armani, May 1979.
- ANL/NDM-49 Some Comments on Resolution and the Analysis and Interpretation of Experimental Results from Differential Neutron Measurements by Donald L. Smith, November 1979.
- ANL/NDM-50 Prompt-Fission-Neutron Spectra of  $^{233}\text{U}$ ,  $^{235}\text{U}$ ,  $^{239}\text{Pu}$  and  $^{240}\text{Pu}$  Relative to that of  $^{252}\text{Cf}$  by A. Smith, P. Guenther, G. Winkler and R. McKnight, September 1979.
- ANL/NDM-51 Measured and Evaluated Neutron Cross Sections of Elemental Bismuth by A. Smith, P. Guenther, D. Smith and J. Whalen, April 1980.
- ANL/NDM-52 Neutron Total and Scattering Cross Sections of  $^6\text{Li}$  in the Few MeV Region by P. Guenther, A. Smith and J. Whalen, February 1980.
- ANL/NDM-53 Neutron Source Investigations in Support of the Cross Section Program at the Argonne Fast-Neutron Generator by James W. Meadows and Donald L. Smith, May 1980.
- ANL/NDM-54 The Nonelastic-Scattering Cross Sections of Elemental Nickel by A. B. Smith, P. T. Guenther and J. F. Whalen, June 1980.
- ANL/NDM-55 Thermal Neutron Calibration of a Tritium Extraction Facility using the  $^6\text{Li}(n,t)^4\text{He}/^{197}\text{Au}(n,\gamma)^{198}\text{Au}$  Cross Section Ratio for Standardization by M. M. Bretscher and D. L. Smith, August 1980.
- ANL/NDM-56 Fast-Neutron Interactions with  $^{182}\text{W}$ ,  $^{184}\text{W}$  and  $^{186}\text{W}$  by P. T. Guenther, A. B. Smith and J. F. Whalen, December 1980.
- ANL/NDM-57 The Total, Elastic- and Inelastic-Scattering Fast-Neutron Cross Sections of Natural Chromium, Peter T. Guenther, Alan B. Smith and James F. Whalen, January 1981.
- ANL/NDM-58 Review of Measurement Techniques for the Neutron Capture Process by W. P. Poenitz, August 1981.
- ANL/NDM-59 Review of the Importance of the Neutron Capture Process in Fission Reactors, Wolfgang P. Poenitz, July 1981.
- ANL/NDM-60 Neutron Capture Activation Cross Sections of  $^{94}\text{Zr}$ ,  $^{96}\text{Zr}$ ,  $^{98,100}\text{Mo}$ , and  $^{110,114,116}\text{Cd}$  at Thermal and 30 keV Energy, John M. Wyrick and Wolfgang P. Poenitz, (to be published).

- ANL/NDM-61 Fast-neutron Total and Scattering Cross Sections of  $^{58}\text{Ni}$  by Carl Budtz-Jørgensen, Peter T. Guenther, Alan B. Smith and James F. Whalen, September 1981.
- ANL/NDM-62 Covariance Matrices and Applications to the Field of Nuclear Data, by Donald L. Smith, November 1981.
- ANL/NDM-63 On Neutron Inelastic-Scattering Cross Sections of  $^{232}\text{Th}$ ,  $^{233}\text{U}$ ,  $^{235}\text{U}$ ,  $^{238}\text{U}$ ,  $^{239}\text{U}$ , and  $^{239}\text{Pu}$  and  $^{240}\text{Pu}$  by Alan B. Smith and Peter T. Guenther, January 1982.
- ANL/NDM-64 The Fission Fragment Angular Distributions and Total Kinetic Energies for  $^{235}\text{U}(n,f)$  from 0.18 to 8.83 MeV by James W. Meadows, and Carl Budtz-Jørgensen, (to be published).
- ANL/NDM-65 Note on the Elastic Scattering of Several MeV Neutrons from Elemental Calcium by Alan B. Smith and Peter T. Guenther, March 1982.
- ANL/NDM-66 Fast-neutron Scattering Cross Sections of Elemental Silver by Alan B. Smith and Peter T. Guenther, May 1982.
- ANL/NDM-67 Non-evaluation Applications for Covariance Matrices by Donald L. Smith, July 1982.
- ANL/NDM-68 Fast-neutron Total and Scattering Cross Sections of  $^{103}\text{Rh}$  by Alan B. Smith, Peter T. Guenther and James F. Whalen (to be published).
- ANL/NDM-69 Fast-neutron Scattering Cross Sections of Elemental Zirconium by Alan B. Smith and Peter T. Guenther (to be published).
- ANL/NDM-70 Fast-neutron Total and Scattering Cross Sections of Niobium by Alan B. Smith, Peter T. Guenther and James F. Whalen (to be published).
- ANL/NDM-71 Fast-neutron Total and Scattering Cross Sections of Elemental Palladium by Alan B. Smith, Peter T. Guenther and James F. Whalen, June 1982.
- ANL/NDM-72 Fast-neutron Scattering from Elemental Cadmium by Alan B. Smith and Peter T. Guenther (to be published).
- ANL/NDM-73 Fast-neutron Elastic-Scattering Cross Sections of Elemental Tin by C. Budtz-Jørgensen, Peter T. Guenther and Alan B. Smith (to be published).
- ANL/NDM-74 Evaluation of the  $^{238}\text{U}$  Neutron Total Cross Section by Wolfgang Poenitz, Alan B. Smith and Robert Howerton (to be published).

## Table of Contents

	<u>Page</u>
ABSTRACT . . . . .	viii
I. INTRODUCTION . . . . .	1
II. EXPERIMENTAL METHODS . . . . .	1
III. EXPERIMENTAL RESULTS . . . . .	2
A. Neutron Total Cross Sections . . . . .	2
B. Neutron Scattering Cross Section . . . . .	2
IV. MODEL DERIVATION AND DISCUSSION . . . . .	6
V. COMPARISONS WITH ENDF/B-V . . . . .	9
VI. SUMMARY COMMENTS . . . . .	9
ACKNOWLEDGEMENTS . . . . .	11
REFERENCES . . . . .	12

NEUTRON TOTAL AND SCATTERING  
CROSS SECTIONS OF ELEMENTAL ANTIMONY\*

by

A. B. Smith, P. T. Guenther and J. F. Whalen

Applied Physics Division  
Argonne National Laboratory  
Argonne, Illinois 60439

ABSTRACT

Neutron total cross sections are measured from 0.8 to 4.5 MeV with broad resolutions. Differential-neutron-elastic-scattering cross sections are measured from 1.5 to 4.0 MeV at intervals of 50 to 200 keV and at scattering angles distributed between 20 and 160 degrees. Lumped-level neutron-inelastic-scattering cross sections are measured over the same angular and energy range. The experimental results are discussed in terms of an optical-statistical model and are compared with respective values given in ENDF/B-V.

---

\*This work supported by the U. S. Department of Energy.

## I. INTRODUCTION

This study was undertaken as part of a series of experimental investigations of the fast-neutron interaction with light-mass fission products. Elemental antimony consists of two essentially equally-abundant isotopes:  $^{121}\text{Sb}$ (57.3%) and  $^{123}\text{Sb}$ (42.7%). Both have small fission-product yields, typically a few times  $10^{-2}\%$ . Thus the isotopes of antimony are not fission products of direct applied interest. However, the element is near the shell closure at  $Z=50$  and at the upper-mass extreme of the light-mass fission yields. Thus the fast-neutron interaction with antimony is of interest in the context of the systematic behavior of the fast-neutron interaction with light-mass fission products. In particular, antimony is a worthwhile reference point for determining model parameters useful for extrapolating observed neutron processes into experimentally inaccessible fission-product regions. The objectives of the present work were the provision of an experimental foundation and the subsequent development of parameters for a relevant optical-statistical model (OM). This and similar models contributed to the formulation of a "regional" OM generally applicable to light-mass fission products, as described elsewhere.<sup>1</sup> Subsequent portions of these remarks deal with the experimental results, the derivation of the model, and comparisons of the present results with values given in ENDF/B-V.<sup>2</sup>

## II. EXPERIMENTAL METHODS

The experimental samples were solid metal cylinders of chemically pure elemental antimony. Those used in the neutron-total-cross-section measurements were 2.5 cm in diameter and 1.25 cm long. They were stacked to obtain neutron transmissions in the range 40% to 60%. The scattering sample was a cylinder 2 cm in diameter and 2 cm long. Sample densities were determined by precise weight and dimension measurements.

The neutron total cross sections were determined from the observed neutron transmissions through the measurement samples in the conventional manner.<sup>3</sup> The neutron source was the  $^7\text{Li}(p;n)^7\text{Be}$  reaction pulsed on for durations of  $\approx 1$  nsec at a repetition rate of 2 MHz. The samples were placed  $\approx 1$  m from the source at a zero-degree reaction angle. The measurement samples and void positions were alternated in the neutron beam  $\approx 24$  times a minute. Neutrons were detected with a proton-recoil scintillator placed 4 to 6 m from the source. Conventional time-of-flight techniques were used to suppress background effects and to select the primary neutron group from the source reaction. In-scattering effects were negligible. A reference pulser was introduced into the measurement system to assure proper corrections for dead-time effects. Neutron total cross sections of carbon were concurrently determined in order to verify the performance of the apparatus.<sup>4</sup> The measurement method and apparatus were identical to those described in detail in ref. 5.

The neutron-scattering measurements utilized the Argonne National Laboratory 10-angle time-of-flight apparatus.<sup>6</sup> The neutron source was, again, the

${}^7\text{Li}(p;n){}^7\text{Be}$  reaction. The neutron detectors were placed at the ends of  $\approx 5.4$  m flight paths distributed over the angular range 20 to 160 degrees. The relative angular scale was determined to within  $\pm 0.2$  degrees and the absolute setting of the angular system to within  $\pm 0.7$  degrees. Below 3.0 MeV the measurements were made at ten angles; at higher energies twenty angles were used. Relative neutron-detector sensitivities were determined by the observation of neutrons emitted from the spontaneous fission of  ${}^{252}\text{Cf}$ .<sup>7</sup> The absolute normalization of these detector efficiencies was determined relative to the neutron total cross sections of carbon<sup>4</sup> in the manner described in ref. 8. This normalization method implies that the antimony scattering cross sections were determined relative to the neutron total cross sections of carbon. All of the measured differential-scattering cross sections were corrected for multiple-event, beam-attenuation and angular-resolution effects as described in ref. 9. The latter reference also contains a detailed description of the measurement apparatus and its applications.

### III. EXPERIMENTAL RESULTS

#### A. Neutron Total Cross Sections

The neutron total-cross-section measurements were made with incident-neutron energy resolutions of 30 to 50 keV from  $\approx 0.8$  to 4.5 MeV at intervals of  $\approx 50$  keV. Several sets of data, covering the entire energy range, were obtained. These sets were averaged over 100 keV intervals to obtain the experimental results shown in Fig. 1. The statistical uncertainties of the averaged values are  $\approx 1\%$ . The results were extended to lower energies (as shown in Fig. 1) using a 50 keV average of the recent experimental values of Poenitz et al.<sup>10</sup> Self-shielding effects were estimated using the methods of ref. 10 and they were found to be negligible in the energy range of the present measurements. The prior neutron-total-cross-section data base should be reasonably summarized by ENDF/B-V<sup>2</sup>. Comparisons with that evaluated data set are discussed below.

#### B. Neutron Scattering Cross Section

The differential scattering measurements were made from  $\approx 1.5$  to 4.0 MeV in steps of  $\approx 50$  keV below 3.0 MeV and steps of  $\approx 200$  keV at higher energies. Energy-adjacent distributions were averaged below 3.0 MeV. The incident-neutron energy spreads were 50 to 70 keV. These relatively large values were chosen so as to average possible energy-dependent fluctuations, enhance statistical accuracies and remove ambiguities due to experimental resolution. The latter were a concern with the elemental antimony targets.  ${}^{121}\text{Sb}$  has a  $5/2+$  ground state and a  $7/2+$  first-excited level at 37 keV while  ${}^{123}\text{Sb}$  has a  $7/2+$  ground state and a  $5/2+$  first-excited level at 160 keV.<sup>12</sup> The complete resolution of these components throughout




Fig. 1. Neutron Total and Elastic-Scattering Cross Sections of Elemental Antimony. The present experimental results are indicated by X and  $\diamond$  (total cross sections extended below 0.8 MeV using 50 keV averages of the recent results of ref. 10). The lower-energy elastic-scattering values of ref. 11 are indicated by O. Curves denote the results of model calculations as described in the text.

the measured energy range would be exceedingly tedious with possibly uncertain results. Therefore, the measurements intentionally included inelastically scattered neutrons due to the excitation of both first-excited levels with the elastically-scattered component. This observational reality was treated in the interpretation, as discussed below.

The differential-elastic-scattering results are summarized in Fig. 2. The distributions behave in an energy-smooth manner. The statistical uncertainties of the individual experimental values were  $\lesssim 1\%$ . The uncertainties associated with the detector calibrations were  $\approx 3\%$ . Correction procedures introduced an additional  $\lesssim 1\%$  uncertainty. Thus the overall differential-elastic-scattering uncertainties were  $\approx 5\%$ . Near the extreme minima of the distributions the statistical uncertainties associated with both the experimental measurements and the correction procedures were larger. The angle-integrated elastic-scattering cross sections were derived by least-square fitting the measured differential values with 6<sup>th</sup>-order Legendre-Polynomial series. The results of these fitting procedures were descriptive of the experimental results, as indicated in Fig. 2. The resulting angle-integrated elastic-scattering cross sections are shown in Fig. 1. Their estimated uncertainties were  $\approx 5\%$  and they are consistent to well within that estimate. These angle-integrated elastic-scattering cross sections are inclusive of inelastic-scattered components due to the excitation of the first-excited levels, as noted above. The present differential and angle-integrated elastic-scattering cross sections are in good agreement with lower-energy results previously reported from this laboratory,<sup>11</sup> as illustrated in Fig. 1. More generally, the prior data base should be reasonably summarized by ENDF/B-V<sup>2</sup> and comparisons of the present experimental elastic-scattering results with that evaluated data set are discussed below.

The broad resolutions employed in the measurements and the complexity of the elemental level structure precluded detailed inelastic-scattering results. However, five lumped-level inelastic excitations were observed. The corresponding excitation energies are given in Table I. These are averages of a number of observations with uncertainties defined as the RMS deviations from the averages. Table I correlates these observed excitations with reported levels in both isotopes.<sup>12</sup> In all cases, the observed inelastic-neutron group is a composite of several components and, in some cases, of many components. Due to this complexity the correlations are only qualitative.

The angle-integrated inelastic-excitation cross sections were derived from the measured differential values by Legendre-Polynomial fitting as described above for elastic scattering. The polynomials were of low order as the observed differential distributions were nearly isotropic. The angle-integrated inelastic-scattering cross-section results are summarized in Fig. 3. The observed excitations of the 491 keV level are uncertain as the magnitudes were much smaller than the cross sections due to the elastic scattering of the second neutron group from the source reaction. Corrections were made for the latter contaminant but at the expense of considerable increases in the experimental uncertainties. Cross sections for the excitation of 1050 and 1436 keV levels are better defined and, in each case,




Fig. 2. Differential Elastic-Scattering Cross Sections of Elemental Antimony. The present experimental results are indicated by data symbols. Curves denote the results of fitting Legendre-Polynomial series to the measured quantities.

must consist of a number of components. The higher-energy excitations are less well known. The inelastic-scattering cross-section uncertainties, shown in Fig. 3, range from 15% to 30%. Their origins are similar to those described above for elastic scattering but with larger statistical components and with contributions due to corrections for the second-neutron component from the source reaction. The present inelastic-scattering results are consistent with the lower-energy values previously reported by this laboratory<sup>11</sup> and with the above neutron total and elastic-scattering cross-section results.

#### IV. MODEL DERIVATION AND DISCUSSION

It was assumed that the present experimental results could be described in terms of a spherical optical-statistical model(OM).<sup>13</sup> The model parameters were obtained by concurrently chi-square fitting the differential-elastic-scattering cross sections of Fig. 2, simultaneously varying the six parameters; real and imaginary strengths, radii and diffusenesses. A real-strength energy dependence of the form  $V=V_0 - 0.3 E(\text{MeV})$  and a spin-orbit strength of 6 MeV were assumed. Compound-nucleus (CN) contributions were calculated using the Hauser-Feshbach formula,<sup>14</sup> with the corrections of Moldauer.<sup>15</sup> The excitation of discrete levels to energies of  $\approx 1.1$  MeV was explicitly calculated using the energies, spins and parities of ref. 12. Higher-energy excitations were described using the statistical formulation and parameters of Gilbert and Cameron.<sup>16</sup>

The application of the above procedures to the particular case of elemental antimony presents two additional problems. The two isotopes have different ground-state spins ( $5/2+$  for  $^{121}\text{Sb}$  and  $7/2+$  for  $^{123}\text{Sb}$ ). As noted above, the present measurements intentionally included neutrons due to the excitation of the first levels with those due to true elastic-scattering. The calculations were carried out using the program ABAREX<sup>17</sup> which has the capability to fit composite distributions such as those measured in this work. In this composite fitting the two isotopes behave similarly as they differ only in alternate ground and first-excited-level configurations; i.e. ( $7/2+$  and  $5/2+$ ) and ( $5/2+$  and  $7/2+$ ). The energies, and therefore the transmission coefficients, are similar. Because of the better known  $^{123}\text{Sb}$  level structure, that isotope was assumed in most of the fitting procedures. The resulting parameters were then used as a starting point for fitting assuming a  $^{121}\text{Sb}$  target. The  $^{121}\text{Sb}$ -based parameters were essentially identical to those obtained with the  $^{123}\text{Sb}$  target. The parameters were further verified by calculating the elemental CE contributions at each energy, subtracting them from the measured values to obtain the "experimental" shape-elastic (SE) cross sections and fitting the latter assuming only SE processes. After several iterations, the SE procedure led to parameters very similar to those obtained using the above  $^{121}\text{Sb}$  and  $^{123}\text{Sb}$  target assumptions. The two parameter sets obtained using the isotopic assumptions were averaged to obtain the elemental parameter set given in Table II. These parameters provide a reasonable description of the observed differential-elastic-scattering cross sections, as illustrated in Fig. 4, and of the neutron total and angle-integrated elastic-scattering cross sections, as illustrated in Fig. 1. However, there are some small discrepancies between measured and calculated results.




Fig. 3. Lumped-level Neutron-Inelastic-Excitation Cross Sections of Elemental Antimony. The present experimental results are indicated by  $\square$ , 200 keV averages of those of ref. 11 by  $+$ . Light curves are "eyeguides" and heavy curves indicate the results of calculations, as described in the text.




Fig. 4. Comparison of Measured and Calculated Differential-Elastic-Scattering Cross Sections of Elemental Antimony. The measured values are indicated by data symbols and the calculated results by curves.

In particular, the calculated differential-elastic-scattering cross sections, in the energy range  $\approx 2.0$  to  $3.0$  MeV, and near the minima of the distributions, are systematically smaller than the measured values. This suggests that the statistical representation of ref. 16 leads to excessive channel competition. This suggestion was not pursued by means of pragmatic statistical-parameter adjustments. Fig. 3 compares some measured and calculated angle-integrated inelastic-scattering cross sections. These comparisons are only qualitative as the correlation of lumped-level cross sections with discrete levels is somewhat uncertain. Despite this shortcoming, the agreement is good, excepting the excitation of the  $1050$  keV level. In the latter case the calculated quantities are systematically smaller than the measured values by amounts that cannot be reasonably attributed to level uncertainties. The discrepancy could again be symptomatic of excessive channel competition due to inappropriate statistical-level parameters. Generally, the OM parameters of Table 2 are consistent with those of a "regional" model applicable in this mass-energy region.<sup>1</sup>

## V. COMPARISONS WITH ENDF/B-V

The ENDF/B-V evaluated-data-file system<sup>2</sup> contains  $^{121}\text{Sb}$  and  $^{123}\text{Sb}$  data sets. These isotopic components were combined to obtain elemental-evaluated-neutron total and elastic-scattering cross sections. The elastic-scattering file was constructed so as to include the first inelastic-scattered component ( $E_x=37$  keV ( $^{121}\text{Sb}$ ) and  $=160$  keV ( $^{123}\text{Sb}$ )) in order to make the evaluated results comparable with the present measured values. The measured and evaluated neutron total and elastic-scattering cross sections are compared in Fig. 5 and some relative numerical comparisons are given in Table III. At high energies the two neutron total cross sections are in good agreement. However, as the energy decreases below  $3.0$  MeV the measured values become increasingly larger than the evaluated quantities with differences amounting to  $6\%$ - $8\%$  at  $0.5$  MeV. Above  $1.5$  MeV the measured elastic-scattering cross sections are smaller than the evaluated quantities by more than  $10\%$  over a rather wide energy range. This difference is approximately twice the experimental uncertainty. The measured neutron total and elastic-scattering cross sections imply nonelastic cross sections that are essentially the total inelastic-scattering cross sections over the energy range of the present measurements (except for the excitation of the two low-lying levels, noted above). The nonelastic cross sections implied by the present measurements are larger than given by the evaluation by amounts that are generally several times the experimental uncertainties. This discrepancy is most serious between  $1.5$  and  $3.0$  MeV. The total inelastic-scattering cross sections are large in this region (e.g.  $2$  b and more).

## VI. SUMMARY COMMENTS

The measurements improve the knowledge of energy-averaged neutron total cross sections of elemental antimony from  $\approx 1.0$  to  $4.5$  MeV. They also




Fig. 5. Comparison of Measured and Evaluated Neutron Total and Elastic-Scattering Cross Sections of Elemental Antimony. "Eyeguides" constructed through the present measured values are indicated by heavy curves. Light curves indicate the respective quantities given by ENDF/B-V.<sup>2</sup> The evaluated elastic-scattering and nonelastic values include the excitation of the first two inelastic-neutron groups with the elastic-scattered component as defined in the text.

provide a good understanding of the effective energy-averaged differential-elastic-scattering cross sections from  $\approx 1.5$  to 4.0 MeV and of the corresponding angle-integrated elastic-scattering cross sections. In addition, the measurements give quantitative information about the broad-resolution inelastic-scattering cross sections. The parameters of a conventional spherical optical-statistical model were deduced from the observed differential-elastic-scattering cross sections. These parameters and the model reasonably described the observables. However, detailed comparisons of measured and calculated cross sections tend to suggest that the conventionally accepted statistical-level parameters are not entirely appropriate.<sup>16</sup> The model parameters are a consideration in the selection of a "regional" model applicable to the light fission-product region, described elsewhere.<sup>1</sup> The experimental results suggest that the ENDF/B-V neutron total cross sections are too small at lower energies and that the evaluated elastic-scattering cross sections are considerably too large in the mid-energy range (e.g. 1.5 to 3.0 MeV). Further, these discrepancies suggest that the evaluated total inelastic-scattering cross sections are too small by significant amounts.

#### ACKNOWLEDGEMENTS

The authors are indebted to Drs. W. Poenitz, P. Moldauer and E. Pennington for helpful comments and the provision of pre-publication data.

REFERENCES

1. A. B. Smith, to be published.
2. Evaluated Nuclear Data File-B, Version-V, Brookhaven National Laboratory Report, ENDF-201, (1979), compiled by R. Kinsey.
3. D. Miller, Fast Neutron Physics, Vol.II, J. Marion and J. Fowler Eds., Interscience Pub., New York (1963).
4. A. Smith, R. Holt and J. Whalen, Argonne National Laboratory Report, ANL/NDM-43 (1978).
5. W. Poenitz, J. Whalen and A. Smith, Nucl. Sci. and Eng., 78 331 (1981).
6. A. Smith, P. Guenther, R. Larsen, C. Nelson, P. Walker and J. Whalen, Nucl. Instr. and Methods, 50 277 (1967).
7. A. Smith, P. Guenther and R. Sjoblom, Nucl. Instr. and Methods, 140 397 (1977).
8. A. Smith and P. Guenther, Argonne National Laboratory Report, ANL/NDM-63 (1982).
9. P. T. Guenther, Elastic and Inelastic Neutron Scattering of Fast Neutrons from the Even Isotopes of Tungsten, Univ. of Illinois Thesis (1977).
10. W. Poenitz and J. Whalen, to be published.
11. A. Smith and R. Hayes, Nucl. Phys., A93 609 (1967).
12. Table of Isotopes, 7th Edition, C. M. Lederer and V. S. Shirley Eds., John Wiley and Sons, Inc., New York (1978).
13. P. Hodgson, Nuclear Reactions and Nuclear Structure, Clarendon Press, Oxford (1971).
14. W. Hauser and H. Feshbach, Phys. Rev., 87 366 (1952).
15. P. A. Moldauer, Phys, Rev., C11 426 (1978); also private communication (1982).
16. A. Gilbert and A. Cameron, Can. Jour. of Phys., 43 1446 (1965).
17. ABAREX, a spherical optical-model program; private communication, P. A. Moldauer (1982).

TABLE I. Observed Inelastic-Neutron Excitations in keV

No.	$E_x(\text{exp})$	Reported Levels <sup>a</sup>	
		$^{121}\text{Sb}$	$^{123}\text{Sb}$
1	491±(?)	508(3/2+) 573(1/2+)	542(3/2+)
2	1050±32 <sup>b</sup>	947(9/2+) 1024(7/2+) 1035(9/2+) 1139(11/2+) 1144(9/2+)	1030(9/2+) 1089(11/2+) 1183(+)
3	1436±33 <sup>b</sup>	1386 1408 1410 1427 1446 1514 1521 1630 1660	1262 1324 1338 1511 1577 1601 1643 1653
4	1805±30 <sup>b</sup>	1740 1820	1730 1740 1810 1860
5	2037±30 <sup>b</sup>	1980 +many	2026 +many

<sup>a</sup>As given in Ref. 12; some additional J- $\pi$  values are given therein.

<sup>b</sup>RMS deviation from the average of a number of measurements.

TABLE II. Optical-Model Parameters for Elemental Antimony

---

Real Potential<sup>a</sup>

Strength $V_0$	=	48.36 MeV
Radius <sup>b</sup> , $r_V$	=	1.245 F
Diffuseness, $a_V$	=	0.620 F
$J_V/A$ <sup>d</sup>	=	429.8 MeV · F <sup>3</sup>

Imaginary Potential<sup>c</sup>

Strength, $W$	=	5.60 MeV
Radius <sup>b</sup> , $r_W$	=	1.210 F
Diffuseness, $a_W$	=	0.627 F
$J_W/A$ <sup>d</sup>	=	53.9 MeV · F <sup>3</sup>

---

<sup>a</sup>Assume saxon form, energy dependence  $V = V_0 - 0.3 \cdot E(\text{MeV})$ , and spin-orbit term of Thomas form with 6 MeV strength.

<sup>b</sup>All radii expressed as  $R = r_i \cdot A^{1/3}$ .

<sup>c</sup>Saxon-derivative form.

<sup>d</sup>Strength expressed as integral per nucleon.

TABLE III. Comparisons of Measured and Evaluated Cross Sections

$E_n$ (MeV)	$\frac{\text{ENDF-EXP}}{\text{ENDF}}$ (in percent)		
	$\sigma_t$	$\sigma_{el}$	$\sigma_{nonel}$
1.0	-6.0	-	-
1.5	-3.5	0	-51
2.0	-3.9	+6	-40
2.5	-2.0	+11	-30
3.0	0.0	+10	-16
3.5	0.0	+11	-13
4.0	-1.6	+11	-11
4.5	-2.5	-	-