

NUCLEAR DATA AND MEASUREMENTS SERIES

ANL/NDM-108

**Some Comments on the Interaction of
Fast-Neutrons with Beryllium**

by

M. Sugimoto, P.T. Guenther, J.E. Lynn, A.B. Smith, and J.F. Whalen

November 1988

**ARGONNE NATIONAL LABORATORY,
ARGONNE, ILLINOIS 60439, U.S.A.**

NUCLEAR DATA AND MEASUREMENTS SERIES

ANL/NDM-108


SOME COMMENTS ON THE INTERACTION OF
FAST-NEUTRONS WITH BERYLLIUM

by

M. Sugimoto, P. T. Guenther, J. E. Lynn,
A. B. Smith and J. F. Whalen

November 1988

108



ARGONNE NATIONAL LABORATORY, ARGONNE, ILLINOIS

Operated by THE UNIVERSITY OF CHICAGO

for the U. S. DEPARTMENT OF ENERGY

under Contract W-31-109-Eng-38

Argonne National Laboratory, with facilities in the states of Illinois and Idaho, is owned by the United States government, and operated by The University of Chicago under the provisions of a contract with the Department of Energy.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ANL/NDM-108

SOME COMMENTS ON THE INTERACTION OF
FAST-NEUTRONS WITH BERYLLIUM*

by

M. Sugimoto,** P. T. Guenther, J. E. Lynn,***
A. B. Smith and J. F. Whalen

November 1988

Keywords:

Neutrons incident on ^9Be . Measured σ_t (1 - 10+ MeV), $d\sigma_{el}/d\Omega$
and $d\sigma_{inel}/d\Omega$ (4.5 - 10 MeV). Reaction mechanisms suggested.

Engineering Physics Division
Argonne National Laboratory
9700 South Cass Avenue
Argonne, Illinois 60439
U. S. A.

* This work supported by the U. S. Department of Energy under Contract W-31-109-Eng-38.

** Visiting scientist from Japan Atomic Energy Institute, Tokai, Japan.

*** Argonne Fellow; permanent address AERE, Harwell, United Kingdom.

NUCLEAR DATA AND MEASUREMENTS SERIES

The Nuclear Data and Measurements Series presents results of studies in the field of microscopic nuclear data. The primary objective is the dissemination of information in the comprehensive form required for nuclear technology applications. This Series is devoted to: a) measured microscopic nuclear parameters, b) experimental techniques and facilities employed in measurements, c) the analysis, correlation and interpretation of nuclear data, and d) the evaluation of nuclear data. Contributions to this Series are reviewed to assure technical competence and, unless otherwise stated, the contents can be formally referenced. This Series does not supplant formal journal publication but it does provide the more extensive information required for technological applications (e.g., tabulated numerical data) in a timely manner.

Copies on microfiche can be obtained by contacting:

NTIS Energy Distribution Center
P.O. Box 1300
Oak Ridge, TN 37831
U.S.A.
Comm. Phone: (615) 576-1301
FTS: 626-1301

INFORMATION ABOUT OTHER ISSUES IN THE ANL/NDM SERIES

A list of titles and authors for reports ANL/NDM-1 through ANL/NDM-50 can be obtained by referring to any report of this Series numbered ANL/NDM-51 through ANL/NDM-76, while those for ANL/NDM-51 through ANL/NDM-76 are listed in any report numbered ANL/NDM-77 through ANL/NDM-101. Requests for a complete list of titles or for copies of previous reports should be directed to:

Section Secretary
Applied Nuclear Physics Section
Engineering Physics Division
Building 316
Argonne National Laboratory
9700 South Cass Avenue
Argonne, Illinois 60439
U.S.A.

ANL/NDM-76 Alan B. Smith and Peter T. Guenther, *Scattering of Fast Neutrons from Elemental Molybdenum*, November 1982.

ANL/NDM-77 Donald L. Smith, *A Least-squares Method for Deriving Reaction Differential Cross Section Information from Measurements Performed in Diverse Neutron Fields*, November 1982.

ANL/NDM-78 A.B. Smith, P.T. Guenther and J.F. Whalen, *Fast-neutron Total and Elastic-scattering Cross Sections of Elemental Indium*, November 1982.

ANL/NDM-79 C. Budtz-Joergensen, P. Guenther, A. Smith and J. Whalen, *Few-MeV Neutrons Incident on Yttrium*, June 1983.

ANL/NDM-80 W.P. Poenitz and J.F. Whalen, *Neutron Total Cross Section Measurements in the Energy Region from 47 keV to 20 MeV*, July 1983.

ANL/NDM-81 D.L. Smith and P.T. Guenther, *Covariances for Neutron Cross Sections Calculated Using a Regional Model Based on Elemental-model Fits to Experimental Data*, November 1983.

ANL/NDM-82 D.L. Smith, *Reaction Differential Cross Sections from the Least-squares Unfolding of Ratio Data Measured in Diverse Neutrons Fields*, January 1984.

ANL/NDM-83 J.W. Meadows, *The Fission Cross Sections of Some Thorium, Uranium, Neptunium and Plutonium Isotopes Relative to ^{235}U* , October 1983.

ANL/NDM-84 W.P. Poenitz and J.W. Meadows, *^{235}U and ^{239}Pu Sample-mass Determinations and Intercomparisons*, November 1983.

ANL/NDM-85 D.L. Smith, J.W. Meadows and I. Kanno, *Measurement of the $^{51}\text{V}(n,p)^{51}\text{Ti}$ Reaction Cross Section from Threshold to 9.3 MeV*, June 1984.

ANL/NDM-86 I. Kanno, J.W. Meadows and D.L. Smith, *Energy-differential Cross-section Measurement for the $^{51}\text{V}(n,\alpha)^{48}\text{Sc}$ Reaction*, July 1984.

ANL/NDM-87 D.L. Smith, J.W. Meadows, M.M. Bretscher and S.A. Cox, *Cross-section Measurement for the $^7\text{Li}(n,n't)^4\text{He}$ Reaction at 14.74 MeV*, September 1984.

ANL/NDM-88 A.B. Smith, D.L. Smith and R.J. Howerton, *An Evaluated Nuclear Data File for Niobium*, March 1985.

ANL/NDM-89 Bernard P. Evain, Donald L. Smith and Paul Lucchese, *Compilation and Evaluation of 14-MeV Neutron-activation Cross Sections for Nuclear Technology Applications: Set I*, April 1985.

ANL/NDM-90 D.L. Smith, J.W. Meadows and P.T. Guenther, *Fast-neutron-spectrum Measurements for the Thick-target $^9\text{Be}(d,n)^{10}\text{B}$ Reaction at $E_d = 7$ MeV*, April 1985.

ANL/NDM-91 A.B. Smith, P.T. Guenther and R.D. Lawson, *On the Energy Dependence of the Optical Model of Neutron Scattering from Niobium*, May 1985.

ANL/NDM-92 Donald L. Smith, *Nuclear Data Uncertainties (Vol.-I): Basic Concepts of Probability*, April 1986.

ANL/NDM-93 D.L. Smith, J.W. Meadows and M.M. Bretscher, *Integral Cross-section Measurements for $^7\text{Li}(n,n't)^4\text{He}$, $^{27}\text{Al}(n,p)^{27}\text{Mg}$, $^{27}\text{Al}(n,\alpha)^{24}\text{Na}$, $^{58}\text{Ni}(n,p)^{58}\text{Co}$ and $^{60}\text{Ni}(n,p)^{60}\text{Co}$ Relative to ^{238}U Neutron Fission in the Thick-target $^9\text{Be}(d,n)^{10}\text{B}$ Spectrum at $E_d = 7$ MeV*, October 1985.

ANL/NDM-94 A.B. Smith, D.L. Smith, P. Rousset, R.D. Lawson and R.J. Howerton, *Evaluated Neutronic Data File for Yttrium*, January 1986.

ANL/NDM-95 Donald L. Smith and James W. Meadows, *A Facility for High-intensity Neutron Irradiations Using Thick-target Sources at the Argonne Fast-neutron Generator*, May 1986.

ANL/NDM-96 M. Sugimoto, A.B. Smith and P.T. Guenther, *Ratio of the Prompt-fission-neutron Spectrum of Plutonium-239 to that of Uranium-235*, September 1986.

ANL/NDM-97 J.W. Meadows, *The Fission Cross Sections of ^{230}Th , ^{232}Th , ^{233}U , ^{234}U , ^{236}U , ^{238}U , ^{237}Np , ^{239}Pu and ^{242}Pu Relative ^{235}U at 14.74 MeV Neutron Energy*, December 1986.

ANL/NDM-98 J.W. Meadows, *The Fission Cross Section Ratios and Error Analysis for Ten Thorium, Uranium, Neptunium and Plutonium Isotopes at 14.74-MeV Neutron Energy*, March 1987.

ANL/NDM-99 Donald L. Smith, *Some Comments on the Effects of Long-range Correlations in Covariance Matrices for Nuclear Data*, March 1987.

ANL/NDM-100 A.B. Smith, P.T. Guenther and R.D. Lawson, *The Energy Dependence of the Optical-model Potential for Fast-neutron Scattering from Bismuth*, May 1987.

ANL/NDM-101 A.B. Smith, P.T. Guenther, J.F. Whalen and R.D. Lawson, *Cobalt: Fast Neutrons and Physical Models*, July 1987.

ANL/NDM-102 D. L. Smith, *Investigation of the Influence of the Neutron Spectrum in Determinations of Integral Neutron Cross-Section Ratios*, November 1987.

ANL/NDM-103 A.B. Smith, P.T. Guenther and B. Micklich, *Spectrum of Neutrons Emitted From a Thick Beryllium Target Bombarded With 7 MeV Deuterons*, January 1988.

ANL/NDM-104 L.P. Geraldo and D.L. Smith, *Some Thoughts on Positive Definiteness in the Consideration of Nuclear Data Covariance Matrices*, January 1988.

ANL/NDM-105 A.B. Smith, D.L. Smith, P.T. Guenther, J.W. Meadows, R.D. Lawson, R.J. Howerton, and T. Djemil, *Neutronic Evaluated Nuclear-Data File for Vanadium*, May 1988.

ANL/NDM-106 A.B. Smith, P.T. Guenther, and R.D. Lawson, *Fast-Neutron Elastic Scattering from Elemental Vanadium*, March 1988.

ANL/NDM-107 P. Guenther, R. Lawson, J. Meadows, M. Sugimoto, A. Smith, D. Smith, and R. Howerton, *An Evaluated Neutronic Data File for Elemental Cobalt*, August 1988.

TABLE OF CONTENTS

	<u>Page</u>
Abstract	ix
I Introduction	1
II Experimental Methods	1
III Experimental Results	2
A. Neutron Total Cross Sections	2
B. Neutron Elastic-Scattering Cross Sections	5
C. Neutron Inelastic Scattering	9
IV Discussion	14
V Summary	18
References	21

SOME COMMENTS ON THE INTERACTION OF
FAST-NEUTRONS WITH BERYLLIUM^{*}

by

M. Sugimoto,^{**} P. T. Guenther, J. E. Lynn,^{***}
A. B. Smith and J. F. Whalen

Argonne National Laboratory
Argonne, Illinois 60439, U. S. A.

ABSTRACT

Neutron total cross sections of elemental beryllium were measured from 1 to > 10 MeV with good precision. Differential neutron elastic-scattering cross sections were measured from 4.5 to 10 MeV at energy intervals of ≈ 0.5 MeV, and for ≈ 100 angular steps distributed between 18° and 160° at each incident energy. Concurrently, differential cross sections for the emission of a discrete inelastic-neutron group corresponding to an excited level at 2.43 ± 0.06 MeV were determined over the same incident-energy and angular range. Angle-integrated elastic-scattering cross sections were deduced from the observed differential values to accuracies of 2.5%, and angle-integrated inelastic-scattering cross sections were established to accuracies of $\leq 10\%$. The experimental results are compared with values given in ENDF/B-V, with attention to discrepancies and implications. Qualitative reaction mechanisms are suggested.

* This work supported by the U. S. Department of Energy under Contract W-31-109-Eng-38.

** Visiting scientist from Japan Atomic Energy Institute, Tokai, Japan.

*** Argonne Fellow, Permanent address, AERE, Harwell, United Kingdom.

I. INTRODUCTION

For many years beryllium has been used as a neutron multiplier in fission-energy systems, and recently it has been a prominent candidate for enhancing the neutron economy of fusion-energy systems.¹ The latter application has been a matter of recent debate, as the results of some integral experiments indicate multiplication considerably less than is predicted by calculations using accepted evaluated basic-data sets.²⁻⁵ Integral tests can identify overall shortcomings in the fundamental underlying data and/or in its calculational use, but they cannot define microscopic-data problems which may be due to errors in the basic cross sections, the neutron-emission spectra, or both. The definition of these specific issues and their resolution is a microscopic data problem. Despite these uncertainties and the strong interest in beryllium as a neutron multiplier in fusion systems, there have been very few careful experimental studies of the scattering and emission of fast neutrons from beryllium. Drake et al.⁶ have measured scattered neutrons and emitted neutron spectra at three energies between ≈ 6 -14 MeV. Hogue et al.⁷ have measured elastic and inelastic scattering cross sections at nine energies between ≈ 7 -15 MeV. There have been several studies near 14 MeV, illustrated by the work of Takahashi et al.⁸ Very little is known between the (n,2n) threshold (≈ 1.85 MeV) and 10 MeV, an energy region which will be fairly well-populated in any fusion-blanket containing appreciable amounts of beryllium, and this energy region certainly is a consideration in the interpretation of data from the usual integral tests. The excited level structure of ⁹Be is reasonably known to about 14 MeV as the result of a number of charged-particle and neutron investigations.⁹ There are low-lying levels of both positive and negative parity. Despite this understanding, the reaction mechanism for neutron interaction with beryllium up to 10-14 MeV is not clear. In particular, the strongly inhibited excitation of low-lying positive parity states through the neutron inelastic-scattering process is not explained.

The present work was undertaken in an effort to improve the experimental understanding of the neutron interaction with beryllium in the 1-10 MeV range, and then to test the validity of evaluated nuclear data files currently used in applications. Concurrently, it was hoped to obtain experimental information that would give some guidance as to the fundamental reaction mechanisms involved. Section II deals briefly with the experimental methods used in the measurements, Section III presents the experimental results, and Section IV discusses them in the context of ENDF/B-V. Section IV also considers the qualitative nature of the reaction mechanism governing the inelastic-scattering processes. A topical summary is given in the final Section.

II. EXPERIMENTAL METHODS

The experimental methods employed in the present neutron total and scattering cross-section measurements have been extensively described elsewhere;¹⁰⁻¹⁴ thus, they are outlined very briefly here.

The measurement samples were cylinders of high-purity elemental beryllium (100% ⁹Be). For the total cross-section measurements, the

cylinders were 2 cm in diameter and 1 cm thick, stacked to produce several thicknesses, thereby giving a range of neutron transmissions with a minimum value of $\approx 25\%$. The neutron beam was incident on the axis of the stacked cylinders. The scattering sample was a cylinder 2 cm in diameter and 2 cm long, with neutrons incident upon the lateral surface. All the cylinders were carefully machined from metal ingots, and the densities calculated from weight and dimension measurements were consistent with those given in the literature.

The neutron source used in the total-cross-section measurements was the ${}^9\text{Be}(d,n){}^{10}\text{B}$ reaction ($E_d = 7.0$ MeV).¹⁵ This reaction provided an intense white source of neutrons extending from below 1 MeV to well above 11 MeV. The scattering measurements used the $\text{D}(d,n){}^3\text{He}$ source, with the deuterium gas contained in a cell ≈ 2.5 cm long at a pressure providing neutron energy spreads at the sample of ≈ 200 keV. The mean neutron energy of the monoenergetic $\text{D}(d,n){}^3\text{He}$ source was determined to $\approx \pm 10$ keV by means of controlling the energy of the incident deuteron beam. Throughout the measurements, the neutron sources were pulsed with burst durations of ≈ 1 nsec. The burst repetition rate in the scattering measurements was 2 MHz, and either that or a lower rate was used in the total cross-section measurements.

The total cross sections were deduced from the measured transmission of the neutrons through the measurement samples.¹⁶ The neutron energy was determined using fast-timing techniques, and a number of sample thicknesses were concurrently studied, using the rotating sample technique described in Ref. 10. A variety of neutron flight paths were used, ranging from ≈ 5 m to ≈ 10 m. Considerable attention was given to background control, and this was verified by concurrent measurements of the well-known carbon total cross section, with results in good agreement with those given in the literature.¹⁷

The neutron scattering measurements were made using the Argonne 10-angle scattering apparatus.^{12-14,18} Measurements were made concurrently at ten scattering angles using flight paths of ≈ 500 cm. The scattered neutrons were observed with liquid-scintillation detectors whose relative energy responses were calibrated using the neutrons emitted from the fission of ${}^{252}\text{Cf}$, as described in Ref. 19. The absolute scales of the ten-detector response functions were determined by observing neutrons scattered from the well-known hydrogen standard.¹⁷ All of the scattering measurements were corrected for multiple-event, beam-attenuation and angular-resolution perturbations using Monte-Carlo procedures.²⁰

III. EXPERIMENTAL RESULTS

A. Neutron Total Cross Sections

In the energy range of the present work (≈ 1 -10 MeV), the beryllium total cross section behaves rather smoothly with energy, with only a single large and broad resonance situated at ≈ 2.7 MeV. In this situation, there is no motivation for seeking very fine experimental resolutions, and, thus, instrument scales were selected so that time resolutions of nominally ≈ 3

nsec were obtained. Combined with the flight paths cited above, these provided energy resolutions considerably better than any known structure in the cross section. The neutron energy scale was determined from the measured neutron flight times and flight paths, and verified by the identification of a number of prominent resonances in the neutron total cross section of carbon.¹⁷ The $^9\text{Be}(d,n)$ source is extremely intense, particularly for neutron energies ≤ 6 MeV.¹⁸ As a consequence, counting statistics were very good up to at least 7 MeV. In order to avoid deadtime distortions due to the high response rates, random signals were inserted into the measurement system in such a manner as to give a reliable measure of deadtimes. The geometry was very good; thus, in-scattering corrections were negligible. In addition, the beryllium total cross sections were measured at selected energies using conventional monoenergetic-source techniques in order to verify the white-source results with a different method where background control is better assured. The measurements were made in a concurrent manner with several sample thicknesses providing transmissions ranging from ≈ 25 -75%. Results obtained with the different sample thicknesses were combined to obtain the results shown in Fig. 1. Since the resolutions were better than the physical structure, self-shielding effects should be negligible. The statistical accuracies associated with the individual datum points range from less than 1% to several percent. Identifiable systematic uncertainties (e.g., sample densities) should be considerably smaller.

Figure 1 shows the comparable cross sections given in ENDF/B-V.²¹ The agreement between the present experimental results and the evaluation is generally very good, even through the ≈ 2.7 MeV resonance area where the comparison is sensitive to the exact energy scales. However, in the ≈ 3.2 -4.5 MeV region, the present results are noticeably larger than the values of ENDF/B-V, by up to ≈ 4 -5%, which is considerably beyond the known experimental uncertainties. It seems that the ENDF/B-V total cross sections are essentially based upon the work of Schwartz et al.²² That set of measurements is very detailed, and the stated uncertainties are small. However, there are two other relatively recent, comprehensive sets of beryllium total cross section data, those of Auchampaugh et al.²³ and Foster and Glasgow.²⁴ These two data sets, in the above-cited 3.2-4.5 MeV region of discrepancy, are in good agreement with one another, and with the present results. Thus, in the 3.2-4.5 MeV region, three reasonably contemporary sets of data are consistent with one another, and they are significantly larger than the experimental values of Ref. 22 or the ENDF/B-V evaluation. In this modified form, the evaluation is believed to be accurate to at least $\approx 1.5\%$ from 1-10 MeV. This uncertainty estimate is supported by an independent comprehensive statistical evaluation by one of the present authors (MS), to be reported elsewhere.²⁵ Interestingly, the latter evaluation tends to follow ENDF/B-V in the 3.2-4.5 MeV region of discrepancy. This happens because the large amount of data (with cited good precisions) from Ref. 22 dominated the statistical evaluation procedures. Such a consequence is illustrative of the problem of using statistical evaluation methods to treat discrepant data sets, particularly where the systematic uncertainties are essentially unknown or are underestimated, as apparently has happened in this case. Judgement remains an inherent part of the evaluation process.




Fig. 1. Neutron total cross sections of beryllium. The present experimental values are noted by "+" symbols. The light curve is ENDF/B-V,²¹ and the heavy curve ENDF/B-V, modified as suggested in the text.

B. Neutron Elastic-Scattering Cross Sections

The elastic-scattering measurements were made at angles distributed between $\approx 18^\circ$ and 160° , from incident energies of 4.5 to 10.0 MeV at increments of ≈ 0.5 MeV. Incident-neutron energy spreads varied from ≈ 200 keV at 4.5 MeV to ≈ 100 keV at 10 MeV. Scattered-neutron energy resolutions were more than sufficient to resolve the elastic-scattered component from inelastically scattered contributions due to the excitation of known states in ^9Be . The observed elastically scattered time-of-flight peak was very prominent at all scattering angles and energies, and it was not noticeably perturbed by any other neutron contributions. The target, ^9Be , is light; therefore, neutrons emitted from the sample after multiple elastic collisions can be significantly degraded in energy. The consequence is a small "tail" on the prominent elastically scattered peaks observed in the measurements. Monte-Carlo correction procedures simulated this perturbation and provided appropriate correction factors. At each measurement period, between twenty and forty differential cross sections were determined at a given incident energy, and there were at least four measurement periods for every incident energy, distributed over several years. The measured results at each incident energy were combined into a single distribution containing ≈ 100 differential cross sections. The minimum number was ≈ 75 differential values, and the maximum over 150. To a very large degree, the results from the different measurement periods are independent; thus, their combination provides a realistic measure of statistical and systematic uncertainties that is not available from single measurements. The only known common systematic factors between the results obtained at different times are associated with the sample density, which was carefully determined to fractional-percent accuracies, and with the Monte-Carlo multiple-scattering correction procedures,²⁰ which have been carefully developed and tested over a number of years. The beryllium measurements were generally made concurrently with other scattering measurements, particularly scattering from carbon. The resulting differential elastic- and inelastic-scattering cross sections of carbon were in good agreement with those reported in the literature,¹⁷ both in cross-section magnitude and scattered-neutron energy scale.

The experimental elastic-scattering results are summarized in Fig. 2. There are over a thousand datum values shown in these distributions. A more detailed illustration of a representative distribution is given in Fig. 3. In this example, one can see that the results obtained from the various measurement periods, four in this case, are consistent within the respective experimental uncertainties. The latter are attributed to several sources. There are statistical uncertainties due to counting the scattered particles, both from the beryllium sample and the hydrogen standard, including foregrounds and backgrounds. They were propagated through the data-reduction procedures and are relatively small (from fractional percent to several percent), as the scattered-neutron intensities were good in these measurements and the elastic-scattering cross sections of beryllium are relatively large at all measurement angles. The uncertainties in the $\text{H}(n,n)$ scattering standard are very small (a percent or less) at the energies of the present experiments. The relative measurement angles were very carefully determined using conventional optical instruments, and their uncertainty has




Fig. 3. A detailed illustration of the elastic scattering of 7.14 MeV neutrons from beryllium. The present experimental values are indicated by "O" symbols, and the curve shows the P_6 fit of Eq. 1 to the measured values. The results are given in the laboratory coordinate system.

a negligible effect on the measured results. The normalizations of the relative angular scales were determined by observing the elastic scattering of neutrons from a heavy target (usually niobium) both left and right of the apparent 0° line, at angles where the respective elastic-scattering cross section of niobium (or similar sample) is rapidly changing with angle. It is believed that this procedure determined the zero point of the angular system to within several tenths of a degree. Moreover, these angular calibrations were repeated at each measurement period, and the variations should be random with time. The effect of angle uncertainty was incorporated in the experimental error estimate but was not large, as the beryllium distributions are not as angular dependent as those of a heavier target.

The Monte-Carlo calculations used to correct the measured values for multiple-event, neutron attenuation and angular-resolution effects were pursued through three iterations with a large number of histories (typically 5×10^6 or 1×10^7), resulting in calculational statistical uncertainties of less than 1%. Moreover, the beryllium elastic distributions are not nearly as anisotropic as those of a heavier target and, thus, the Monte-Carlo corrections are more easily carried out to good statistical accuracy. As in all observations, there may be unidentified sources of uncertainty. It was assumed that the latter unidentified contributions amounted to 1.5 mb/sr. The neutrons scattered from the beryllium and the hydrogen standard appear at different energies. Accordingly, the results depend upon the knowledge of the relative energy sensitivities of the individual detectors. The uncertainties in these relative sensitivities are difficult to quantify but are believed to be small (on the order of several percent), and they should be randomly distributed from one measurement to the next. The various uncertainty components were combined in quadrature. Finally, it was subjectively assumed that the minimum overall uncertainty associated with a differential value is $\approx 3\%$ (reflecting possible unidentified systematic uncertainties associated with the detector calibrations), and that became the maximum uncertainty limit in many cases. The measured values are consistent with these uncertainty estimates, as shown in Fig. 3, and one should again remember that this typical example consists of the accumulation of data from four independent measurement periods.

The angle-integrated elastic-scattering cross sections were deduced from the present measurements by least-square fitting the measured values with Legendre-polynomial expansions given by

$$\frac{d\sigma}{d\Omega} = \frac{\sigma}{4\pi} \left[1 + \sum_{i=1}^n \omega_i P_i \right] , \quad (1)$$

where P_i is the i^{th} Legendre polynomial, and n was taken to be 6 or 8. The results obtained with n equal to 6 or 8 agreed to within better than 1%. The value $n = 8$ was used to obtain the cross sections shown in Fig. 4. The uncertainties associated with the angle-integrated elastic-scattering cross sections were believed to be $\approx 2.5\%$. That value was arrived at by fitting




Fig. 4. Comparison of the angle-integrated elastic-scattering results implied by the present work (●) and by the measurements of Hogue et al. (X),⁷ with the values given in ENDF/B-V (curve).

the experimental results of each measurement period with Eq. 1 and comparing the resulting angle-integrated cross section values. There were enough independent measurement periods to give a reasonable sample of the uncertainty in the angle-integrated values. Uncertainties obtained from the least-squares fitting procedure alone were generally smaller; thus, the 2.5% estimate is believed to be conservative.

There are some other comparable elastic-scattering measurements reported in the literature. Perhaps the better and more comprehensive of these is the work by Hogue et al.⁷ These authors report four values in the present measurement range, and their work extends to higher energies (to ≈ 15 MeV). Hogue et al. give their experimental results as coefficients of a Legendre-polynomial expansion in the center-of-mass system. When the present results are converted to the center-of-mass system and refitted with Legendre-polynomial expansions, coefficients very similar to those given by Hogue et al. are obtained. Furthermore, the graphical distributions given by Hogue et al. look very similar to the present experimental results. The resulting angle-integrated elastic-scattering results are compared in Fig. 4. The agreement between the present work and that of Hogue et al. is remarkably good. All of the values are consistent within the respective uncertainties, and for three of the four comparable energies, the agreement is nearly exact (far better than should be expected from the respective uncertainties). Drake et al.⁶ have reported elastic scattering results at ≈ 6 , 10, and 14 MeV. Their angle-integrated elastic-scattering results are in very good agreement with those of the present work, but their uncertainties exceed 10%. Figure 4 also compares the present results, and those of Hogue et al., with the elastic-scattering cross section given in the evaluation of ENDF/B-V.²¹ The measured and evaluated results are somewhat different in both shape and magnitude. The implication of these differences will be discussed in Sec. IV, below.

C. Neutron Inelastic Scattering

There are seven levels in ${}^9\text{Be}$ that are energetically available for inelastic scattering at the energies of the present experiments;⁹ 1.69 MeV ($1/2^+$), 2.429 MeV ($5/2^-$), 2.8 MeV ($1/2^-$), 3.05 MeV ($5/2^+$), 4.70 MeV ($3/2^+$), 6.76 MeV ($7/2^-$), and 7.94 MeV ($1/2^-$). The latter three were not examined, as the respective levels are broad and the resulting inelastically-scattered neutrons have relatively low energies. As a consequence, the emitted neutrons are lost in a continuum spectrum of neutrons emitted in (n,2n) and other breakup reactions. The observed time-of-flight spectra in the region of the first four levels consisted of a broad "hump" of emitted neutrons on which was superimposed a well-resolved and prominent discrete inelastic-neutron group corresponding to an observed excitation of 2.43 ± 0.06 MeV. The underlying "hump" was angular dependent, and, at higher energy, was peaked in the forward direction. At all but the most forward scattering angles, the experimental resolution was sufficient to observe neutrons resulting from the inelastic excitation of the 1.69 MeV level, even given its rather broad width of $\Gamma \approx 210$ keV.⁹ Early photo-emulsion studies²⁶ at incident energies of ≈ 3 MeV qualitatively indicate some inelastic neutrons due to the excitation of the 1.69 MeV level, but there was no evidence for

such a contribution in the present measurements. If such a component is present, it must have a very small cross section. The second level at 2.429 MeV is reported⁹ to have a rather narrow width and closely corresponds to the very prominent inelastic-scattering group observed in the present measurements. The 2.8 MeV level is very broad⁹ ($\Gamma \approx 1.1$ MeV). In the present experiments, it would be impossible to distinguish any corresponding inelastically scattered neutrons from the general broad continuum distribution cited above. The 3.05 MeV level is also comparatively broad ($\Gamma \approx 300$ keV), and is centered only about 600 keV above the 2.429 level. This relatively close spacing and broad width make it difficult to resolve it from the contribution due to the prominent 2.429 MeV level. Inelastic excitation of the 3.05 MeV has been reported,⁶ but at intensities much below that due to the 2.429 MeV level. In the present work there was some indication of a weak inelastic contribution due to the excitation of the 3.06 MeV level, particularly at the larger scattering angles. However, it was of very low intensity and not well defined.

In view of the above situation, the present measurements gave emphasis to determining the inelastic-scattering cross section due to the excitation of the 2.429 MeV level. It was assumed that the continuum emission contribution underlying the prominent experimental peak, including possible artifacts due to the 3.06 MeV level, could be subtracted by linearly extrapolating between the two shoulders of the prominent peak. Subjective judgements were involved in the determination of the shoulder points to be used in the extrapolation, particularly with respect to the lower-energy side of the peak. At different times, these judgements were independently made by two of the authors (MS and AS), with results that were relatively consistent. The objective was always to obtain the contribution attributable to the 2.429 MeV level as free from other components as possible.

The experimental inelastic-scattering results, obtained using the above procedures, are summarized in Fig. 5, and a detailed drawing of one of these distributions is shown in Fig. 6. Each distribution contained a minimum of approximately forty points, and the maximum number exceeded one hundred. Each illustrated distribution is the combination of the results of three or more measurements taken over a several-year period with, as for the elastic-scattering cited above, different instrumental arrangements. The results obtained during the different measurement periods should be essentially independent, and, thus, reproducibility is an indication of uncertainty. The uncertainties attributed to the individual datum points varied a great deal, depending upon the particular energy, angle, and experimental conditions, with the components outlined above in the context of elastic scattering. However, a minimum systematic uncertainty of 10% was assigned to all the measured values as a subjective estimate of uncertainties involved in the above-outlined continuum-subtraction and -summing procedures. At large scattering angles and lower energies, the experimental uncertainties can considerably exceed this minimum value. This increase is due to counting statistics and to the same minimum uncertainty of 1.5 mb/sr, as accepted for the elastic-scattering measurements. As is illustrated in Fig. 6, the data are consistent to well within the uncertainty estimates.




Fig. 5. Neutron differential cross sections for the excitation of the 2.429 MeV level in ${}^9\text{Be}$. The present measured values are indicated by "O" symbols, and the results of fitting Eq. 1 to the data are represented by the curves. The data are given in the laboratory coordinate system.




Fig. 6. Differential cross sections for the excitation of the 2.429 MeV level in ${}^9\text{Be}$ by 7.5 MeV incident neutrons. The measured values are indicated by "O", and the result of a fit of Eq. 1 to the data is represented by the curve. The quantities are given in the laboratory coordinate system.

The angle-integrated cross sections for the excitation of the 2.429 MeV level were deduced from the measured differential values by again least-square fitting Eq. 1 to the measured differential quantities, using a value of $n = 6$. The results of the fitting were descriptive of the measured values, as shown by the curves in Fig. 5. The deduced angle-integrated values are shown in Fig. 7. Their estimated uncertainties are 10%, reflecting the above-mentioned subjective estimates used in the summing procedures. Statistical and other error sources are generally much smaller, and the angle-integrated values obtained from results of different measurement periods are generally consistent to well within the 10% estimate. Figure 7 also shows the results reported by Hogue et al.⁷ and by Drake et al.⁸ The latter values are somewhat larger than the present results, or those of Ref. 7, but have uncertainties of 20-30%. The results of Hogue et al. are systematically larger than the present results, but are only slightly beyond the respective experimental uncertainties, if that. The results of Ref. 7 explicitly include any possible contribution from the 3.06 and 1.69 MeV levels. Therefore, the cross sections should be somewhat larger than those of the present work (by an amount reflecting the two possible additional contributions), which sought to define the inelastic-scattering due to the excitation of the 2.429 MeV level alone. It is interesting that the energy-dependent shapes of the present data and those of Hogue et al. are very similar, even to the small discontinuity in the region of 10 MeV. Figure 7 also shows the comparable inelastic cross section as given in ENDF/B-V.²¹ It is much larger than the present measured values, or those of Hogue et al., by a factor of up to nearly two. This discrepancy is discussed in Sec. IV, below.

IV. DISCUSSION

The present total cross section measurements, and an evaluation by one of the authors (MS),²⁵ support the ENDF/B-V²¹ total cross sections over a very large portion of the region of the present scattering measurements (≈ 5 -10 MeV). The evaluation, and a detailed inspection of the general total cross section data base, indicate that the uncertainties to be associated with the ENDF/B-V total cross sections are, conservatively, $\approx 1.5\%$ or less from ≈ 5 -10 MeV. Therefore, the ENDF/B-V total cross sections, with uncertainties of 1.5%, were assumed for the following discussion. The numerical values at the energies of the present scattering measurements are given in Table 1. The same table gives the corresponding angle-integrated elastic-scattering cross sections deduced by Legendre-polynomial fitting the present data, as described in Sec. III-B. The RMS difference between the measured elastic-scattering cross sections and those given in ENDF/B-V is $\approx 4\%$, with a systematic trend for the evaluated quantities to be 3-5% larger in the middle of the energy range, and nearly the same amount smaller at the upper and lower extremes of the energy range. The differences, while not large, are probably real, as they are approximately twice the experimental uncertainties and are systematic with energy. The nonelastic cross sections ($\sigma_{\text{nonel}} = \sigma_{\text{t}} - \sigma_{\text{el}}$) implied by the measured elastic-scattering cross sections and the total cross sections are also given in Table 1. Of course, they are different from those implied by ENDF/B-V by amounts reflecting the above-cited differences in the elastic-scattering cross sections.




Fig. 7. Angle-integrated cross sections for the excitation of the 2.429 MeV level in ${}^9\text{Be}$. The presently measured values are indicated by "O" symbols, those of Ref. 6 by "+", and of Ref. 7 by "X". The curve indicates the comparable quantities given in ENDF/B-V.²¹

Table 1. Comparative Measured and Evaluated Neutron Cross Sections of Beryllium

Reaction	Incident Neutron Energy in MeV											
	4.5	5.0	5.5	5.9	6.5	7.14	7.5	8.03	8.4	9.06	9.5	10.0
$\sigma_T(b)^{a,b}$	1.9454 (1.5%)	1.8888 (1.5%)	1.8680 (1.5%)	1.8526 (1.5%)	1.8286 (1.5%)	1.7969 (1.5%)	1.7757 (1.5%)	1.7532 (1.5%)	1.7361 (1.5%)	1.7114 (1.5%)	1.6882 (1.5%)	1.6651 (1.5%)
$\sigma_{el}(b)$ Exp. ^c	1.370 (2.5%)	1.300 (2.5%)	1.215 (2.5%)	1.200 (2.5%)	1.150 (2.5%)	1.120 (2.5%)	1.100 (2.5%)	1.145 (2.5%)	1.155 (2.5%)	1.150 (2.5%)	1.145 (2.5%)	1.140 (2.5%)
ENDF/B-V ^d	1.330	1.267	1.240	1.224	1.205	1.186	1.170	1.150	1.138	1.120	1.100	1.080
$\sigma_{nonel}(b)^e$ Exp. ^c	0.5754 (7.8%)	0.5886 (7.3%)	0.6530 (6.3%)	0.6526 (6.3%)	0.6786 (5.9%)	0.6769 (5.8%)	0.6757 (5.6%)	0.6082 (6.4%)	0.5811 (6.7%)	0.5614 (6.9%)	0.5432 (7.0%)	0.5251 (7.2%)
ENDF/B-V	0.6154	0.6219	0.6280	0.6286	0.6236	0.6109	0.6057	0.6032	0.5981	0.5914	0.5882	0.5851
$\sigma_{inel}(b)^f$ Exp. ^c	0.260 (10%)	0.256 (10%)	0.278 (10%)	0.263 (10%)	0.260 (10%)	0.260 (10%)	0.235 (10%)	0.224 (10%)	0.218 (10%)	0.184 (10%)	0.164 (10%)	0.197 (10%)
ENDF/B-V	0.316	0.354	0.389	0.413	0.443	0.450	0.450	0.445	0.434	0.392	0.384	0.375
$\sigma_{nonel} - \sigma_{inel}(b)^g$ Exp. ^c	0.3154 (16%)	0.3326 (15%)	0.3750 (13%)	0.3896 (12%)	0.4186 (11%)	0.4169 (11%)	0.4407 (11%)	0.3842 (12%)	0.3631 (12%)	0.3774 (11%)	0.3792 (11%)	0.3281 (13%)
ENDF/B-V	0.2994	0.2678	0.2386	0.2156	0.1806	0.1609	0.1557	0.1582	0.1640	0.1993	0.2046	0.2101
% Difference ^h	(-5.3%)	(-24%)	(-57%)	(-81%)	(-131%)	(-159%)	(-183%)	(-143%)	(-121%)	(-89%)	(-85%)	(-56%)

16

^aTaken from ENDF/B-V assuming 1.5% uncertainty. See text for details.

^cMeasurement cross sections determined in present experiments, as relevant.

$$^e \sigma_{nonel} = \sigma_T - \sigma_{el}$$

^g $\sigma_{nonel} - \sigma_{inel}(E_x = 2.43 \text{ MeV})$ = remaining partial cross sections, (n,2n), etc.

^bUncertainties in percent are given in brackets.

^dElastic cross sections of ENDF/B-V.

^fInelastic cross section for excitation of 2.43 MeV level only.

^hDefined as (ENDF - EXP)/ENDF in percent.

Table 1 gives the inelastic-scattering cross sections for the excitation of the 2.429 MeV level (σ_{inel}) deduced from the present measurements, as described in Sec. III-C. The respective uncertainties are dominated by the systematic 10% uncertainties conservatively attributed to the summing procedures outlined in Sec. III-C. The table also gives what seem to be the comparable values from ENDF/B-V. The latter are larger than those deduced from the measured values by factors ranging from ≈ 1.2 to more than 2. These are large differences in cross sections that are a substantive fraction of the total nonelastic cross section, and well beyond the uncertainties of the present experimental results. The situation does not substantively change if one considers the alternate experimental results of Hogue et al.⁷ These differences are also clearly evident in Fig. 7. The experimental measurements also imply the totality of partial nonelastic cross sections, exclusive of the inelastic cross sections for the explicit excitation of the 2.429 MeV level ($\sigma_{\text{part}} \equiv \sigma_{\text{nonel}} - \sigma_{\text{inel}}$). Those partial cross sections include a number of reaction channels, in particular contributions to the (n,2n) reactions. The corresponding experimentally deduced quantities are given in Table 1. The uncertainties are in the range 11-16%. The comparable ENDF/B-V values systematically vary from reasonable agreement at the lowest energies ($\approx 5\%$, the low-energy value must be taken with caution due to the σ_t discrepancy noted above) to very large differences at the higher energies, with the experimentally deduced values more than a factor of 2.5 larger at some energies, and, generally, approximately a factor of 2 larger over a major portion of the energy range. Moreover, the experimentally deduced cross sections are large, $\approx 20\%$ of the total cross section. Clearly, the present measurements suggest some rather significant discrepancies with the ENDF/B-V partial cross sections, and these may be reflected in integral energy-transfer mechanisms. The implication of these differences may be considerable in some applications, but the details will depend on the exact nature of the other partial cross sections, particularly the continuum neutron-emission processes.

As noted in Section III-C, above, the inelastic-neutron excitation of the first excited state at 1.69 MeV ($1/2^+$) is very weak and was not observed in the present measurements. The next positive parity state (3.05 MeV, $5/2^+$) is also very weakly excited, while, at the same time, scattering to the odd parity states is prominent (ground state, $3/2^-$, and 2.429 MeV, $5/2^-$, level). This suggests a possible shape "co-existence" in the ${}^9\text{Be}$ nucleus. The Nilsson diagram for the lightest nuclei suggests that the last neutron in the ground-state ($J^\pi = 3/2^-$) configuration of ${}^9\text{Be}$ should lie in the $[101\ 3/2^-]$ orbit (stemming from the $1p_{3/2}$ state of the spherical potential), and, to be energetically favorable, should tend to push the nucleus towards a spherical shape. The first-excited state ($J^\pi = 1/2^+$) should, on the other hand, have its final neutron in the $[220\ 1/2^+]$ orbital (from the $1d_{5/2}$ state), which, to favor energy, will tend to push the nuclear shape towards large prolate

deformation. The very low excitation of this state (1.69 MeV) certainly favors this interpretation, and its large width ($\Gamma \approx 200$ keV) also indicates the large deformation preceding breakup into the two α -particles underlying the ${}^9\text{Be}$ core structure. Most of the higher states appear to be interpretable as rotational bands, with these first two states as band heads. Thus, the second excited state (2.429 MeV, $5/2^-$) appears to be the first rotational level built upon the ground state, while the $5/2^+$ state at 3.05 MeV, and probably the $3/2^+$ state at 4.7 MeV, could be members of a decoupled band based upon the 1.69 MeV state. If such interpretations are correct, the rotational moment-of-inertia parameters of the bands are, respectively, $\hbar^2/2I \approx 0.5$ MeV and 0.34 MeV, again indicating a much greater deformation for the 1.69 MeV state than for the ground state.

The interaction of an incident neutron in the zero to several-MeV range with the ${}^9\text{Be}$ ground state will develop through states that can be described as a single-particle neutron orbiting in the field of the near-spherical ${}^9\text{Be}$ $3/2^-$ core. The relevant single-particle orbits will be those Nilsson orbits developing from the spherical $2s_{1/2}$ and $1d_{3/2}$ states, and, to a much less extent, from the $1d_{5/2}$ state. The predominant total J^π values concerned are 1^- and 2^- , to a lesser extent 0^- and 3^- , and to a small degree 4^- . These states have little similarity to the configurations required to give a significant reduced neutron width for emission to the first-excited state of ${}^9\text{Be}$. For $J^\pi = 0^-$, 1^- , and 2^- states, the $[101\ 3/2]$ and $[101\ 1/2]$ orbitals (both lying much lower than the s and d orbitals) are required, and for the $J^\pi = 0^-$ and 1^- states alone, the $[330\ 1/2]$ orbit stemming from the $1f_{7/2}$ spherical state will be involved. Because of the large deformation expected for the ${}^9\text{Be}$ excited core, this last orbit can be expected to lie in the same energy region as the 2s and 1d orbitals. This alone could favor relatively strong mixing of "initial" and "final" configurations. However, against such mixing, one could expect the rather large difference in core deformation to provide a decisive hindrance factor. Thus, one expects the states strongly excited by neutron absorption in ${}^9\text{Be}$ to have weak reduced widths for neutron reemission to the first excited state. In contrast, the configurations involved in neutron emission to the second excited state are expected to be mixed strongly, through rotational coupling, with those invoked in the initial neutron absorption mechanism.

V. SUMMARY

The present neutron total-cross-section measurements support the ENDF/B-V^{2.1} evaluation from 1-10+ MeV, excepting a local region between approximately 3.2 and 4.5 MeV where the present experimental values, and those in the literature, suggest that the evaluated total cross sections should be increased by up to $\approx 5\%$. The present measurements give new definition to the differential and angle-integrated elastic- and inelastic-

scattering cross sections over the incident energy range 4.5-10 MeV. The experimental elastic-scattering results differ modestly from those given in ENDF/B-V. They have been used in upgrading the evaluated elastic-scattering cross sections of the proposed ENDF/B-VI, as illustrated by the comparisons of Fig. 8.²⁷ The cross sections for the inelastic-neutron excitation of the 2.429 MeV level were determined with results that disagree with those apparently given in ENDF/B-V by up to a factor of approximately two. These are large differences in a cross section that is a significant part of the total cross section. The experimental results imply that the remainder of the nonelastic cross section, exclusive of the excitation of the 2.429 MeV level, are very different from those given in ENDF/B-V (by a factor of two or more) over the large majority of the the 4.5-10 MeV measured energy range. Such differences can be a serious concern in a diversity of applications to fusion- and fission-energy systems which rely upon neutron multiplication in beryllium for viability. The details of this concern depend upon the character of the various partial cross sections and the associated emission spectra, which are not well known. The quantitative nature of the nuclear processes governing the interaction of fast neutrons with ⁹Be are not well understood. It is suggested that, qualitatively, the processes leading to the excitation of the ground and low-lying states are governed by two rotational bands based upon the negative-parity ground state and the positive-parity first-excited (1.69 MeV) state, in a manner that strongly inhibits inelastic-neutron excitation of the positive-parity levels. The concepts are consistent with the observed results, and with the collective model of Nilsson.

The interaction of fast-neutrons with ⁹Be remains somewhat of an enigma from both fundamental and applied points of view. Integral studies are useful to test the overall processes, but they will not provide sufficient information to resolve the microscopic physical questions raised above. They are necessary but not sufficient criteria for testing or validating the underlying physical processes. What is needed for applied and fundamental understanding are detailed microscopic experimental and theoretical studies directed toward a comprehensive understanding of the interaction of fast neutrons with ⁹Be. In particular, these should include precise studies of discrete neutron inelastic scattering and of continuum neutron-emission processes using the best contemporary techniques, extending well beyond the limited objectives of the present work and the information available in the literature. It is indeed remarkable that so little reliable neutron information is available for a nucleus of such interest for important applications.




Fig. 8. Illustrative use of the present experimental results to upgrade the proposed ENDF/B-VI ${}^9\text{Be}$ evaluation.²⁷ The present elastic-scattering results are indicated by vertical bars, the evaluation by the curves. Incident-neutron energies are 5.5 and 8.4 MeV. The data shown are in the center-of-mass system.

REFERENCES

1. E. T. Cheng, private communication (1988).
2. R. W. Moir, private communication (1988).
3. T. K. Basu, V. R. Nargundkar, P. Cloth, D. Filges and S. Taczanowski, Nucl. Sci. and Eng. 70 309 (1979); see also Fusion Energy 12 381 (1987).
4. V. A. Zagryadskij et al., IAEA Report, INDC(CCCP)-272/G (1987).
5. R. S. Hartly, Ph.D. dissertation, University of Texas (1987).
6. D. Drake, G. Auchampaugh, E. Arthur, C. Ragan and P. Young, Nucl. Sci. and Eng. 63 401 (1977).
7. H. Hogue, P. Von Behren, D. Epperson, S. Glendinning, P. Lisowski, C. Nelson, H. Newson, F. Purser, W. Tornow, C. Gould and L. Seagondollar, Nucl. Sci. and Eng. 68 38 (1978).
8. A. Takahashi et al., Japan Atomic Energy Research Institute Report, JAERI-M-88-065, p. 279 (1988).
9. F. Ajzenberg-Selove, Nucl. Phys. A413 1 (1984).
10. W. P. Poenitz, A. B. Smith and J. F. Whalen, Nucl. Sci. and Eng. 78 333 (1981).
11. W. Poenitz and J. Whalen, Argonne National Laboratory Report, ANL/NDM-80 (1983).
12. C. Budtz-Jorgensen, P. Guenther, A. Smith, J. Whalen, W. McMurray, M. Renan and I. Van Heerden, Z. Phys. A319 47 (1984).
13. A. Smith, P. Guenther, J. Whalen, I. van Heerden and W. McMurray, J. Phys. G11 125 (1985).
14. P. Guenther, A. Smith and J. Whalen, Phys. Rev. C12 1797 (1976).
15. A. Smith, P. Guenther and B. Micklich, Argonne National Laboratory Report, ANL/NDM-103 (1988).
16. D. Miller, Fast Neutron Physics Vol.-II. Ed. J. Marion and J. Fowler, Interscience Publishers, New York (1963).
17. Nuclear Data Standards for Nuclear Measurements, IAEA Tech. Report No. 227, Editors H. Conde, A. Smith and A. Lorenz, IAEA Press (1983).
18. A. Smith, P. Guenther, R. Larson, C. Nelson, P. Walker and J. Whalen, Nucl. Instr. Methods 50 277 (1967).
19. A. Smith, P. Guenther and R. Sjoblum, Nucl. Instr. Methods 140 397 (1977).

20. MONTE-SPHERE and MONTE-POLY. Monte-Carlo computational codes, A. Smith, unpublished memorandum (1988). Codes available from the author.
21. Evaluated Nuclear Data File-B, Version-V, National Nuclear Data Center, Brookhaven National Laboratory.
22. R. Schwartz, R. Schrack and H. Heaton, National Bureau of Standards Monograph, NBS-138 (1974).
23. G. Auchampaugh, S. Plattard and N. Hill, Nucl. Sci. and Eng. 69 30 (1979).
24. D. Foster and D. Glasgow, Phys. Rev. C3 576 (1971).
25. M. Sugimoto et al., Argonne National Laboratory Report (to be published).
26. R. Wagner and P. Huber, Helv. Phys. Acta 23 89 (1957).
27. R. J. Howerton, private communication (1988).